

Stability index of linear random dynamical systems*

Anna Cima⁽¹⁾, Armengol Gasull⁽¹⁾ and Víctor Mañosa⁽²⁾

⁽¹⁾ *Departament de Matemàtiques, Facultat de Ciències,
Universitat Autònoma de Barcelona,
08193 Bellaterra, Barcelona, Spain
cima@mat.uab.cat, gasull@mat.uab.cat*

⁽²⁾ *Departament de Matemàtiques,
Universitat Politècnica de Catalunya
Colom 1, 08222 Terrassa, Spain
victor.manosa@upc.edu*

April 11, 2019

Abstract

Given a homogeneous linear discrete or continuous dynamical system, its stability index is given by the dimension of the stable manifold of the zero solution. In particular, for the n dimensional case, the zero solution is globally asymptotically stable if and only if this stability index is n . Fixed n , let $p_k, k = 0, 1, \dots, n$, denote the probabilities that the random variable that assigns to each linear random dynamical system its stability index takes the value k . In this paper we obtain either the exact values p_k , or their estimations by combining the Monte Carlo method with a least square approach that uses some affine relations among the values $p_k, k = 0, 1, \dots, n$. The particular case of n -order homogeneous linear random differential or difference equations is also studied in detail.

Mathematics Subject Classification 2010: 37H10, 34F05, 39A25, 37C75.

Keywords: Stability index; random differential equations; random difference equations; random dynamical systems.

*The authors are supported by Ministry of Economy, Industry and Competitiveness–State Research Agency of the Spanish Government through grants MTM2016-77278-P (MINECO/AEI/FEDER, UE, first and second authors) and DPI2016-77407-P (MINECO/AEI/FEDER, UE, third author). The first and second authors are also supported by the grant 2017-SGR-1617 from AGAUR, Generalitat de Catalunya. The third author acknowledges the group’s research recognition 2017-SGR-388 from AGAUR, Generalitat de Catalunya. We want to acknowledge Prof. Maria Jolis for her helpful indications.

1 Introduction

Nowadays it is unnecessary to emphasize the importance of ordinary differential equations and discrete dynamical systems to model real world phenomena, from physics to biology, from economics to sociology. These dynamical systems, a concept that includes both continuous and discrete models (and even dynamic equations in time-scales), can have undetermined coefficients that in the case of real applications, must be adjusted to fixed values that serve to make good predictions: this is known as the identification process. Once these coefficients are fixed we obtain a deterministic model.

In recent years some authors have highlighted the utility of considering random rather than deterministic coefficients to incorporate effects due to errors in the identification process, natural variability in some of the physical parameters, or as a method to treat and to incorporate uncertainties in the model, see [4, 5, 16] for examples coming from biological modeling and [9] for examples coming from mechanical systems.

In the same aim that inspires some works like [1, 11], in this paper we focus on giving a statistical measure of the stability for both discrete and continuous linear dynamical systems,

$$\dot{\mathbf{x}} = A \mathbf{x} \quad \text{or} \quad \mathbf{x}_{k+1} = A \mathbf{x}_k, \quad (1)$$

where both $\mathbf{x}, \mathbf{x}_k \in \mathbb{R}^n$ and A is an $n \times n$ real matrix.

More concretely, in the continuous (resp. discrete) case we define the *stability index* of the origin, $s(A)$, as the number of eigenvalues, taking into account their multiplicities, of A with negative real part (resp. modulus smaller than 1). This index coincides with the dimension of the invariant stable manifold of the origin. Notice also that if $s(A) = n$ (resp. $s(A) = 0$) the origin is a global stable attractor (resp. a global unstable repeller).

In this work we study the probabilities p_k for a linear dynamic system (1) to have a given stability index k when the parameters of the matrix A are random variables with a given natural distribution. As we will see in Section 2, this distribution must be that all the elements of A are *independent and identically distributed* (i.i.d.) normal random variables with zero mean. We also will study the same question for linear n -th order differential equations and for linear difference equations.

We also remark that our results can be extrapolated to know a measure of the stability behaviour of critical or fixed points for general non-linear dynamical systems, because near them they can be written as

$$\dot{\mathbf{x}} = A \mathbf{x} + f(\mathbf{x}), \quad \text{or} \quad \mathbf{x}_{k+1} = A \mathbf{x}_k + f(\mathbf{x}_k),$$

with f being a non-linear term vanishing at zero. Moreover, while the situation where the origin is non-hyperbolic is negligible, in the complementary one, the stability index of the

linear part coincides with the dimension of the local stable manifold at the point.

The key tool in the continuous case to know the stability index of a matrix is the Routh-Hurwitz criterion, see for instance [8, p. 1076]. This approach allows to know the number of roots of a polynomial with negative real part in terms of algebraic inequalities among its coefficients. Similarly, its counterpart for the discrete case is called Jury criterion. It is worth to observe that in fact both are equivalent and it is possible to get one from the other by using a Möbius transformation that sends the left hand part of the complex plane into the complex ball of radius 1.

In all the cases, when we do not know how to compute analytically the searched probabilities, we introduce a two steps approach to obtain estimations of them:

- **Step 1:** We start using the celebrated Monte Carlo method. Recall that this computational algorithm relies on repeated random sampling and gives estimations of the searched probabilities based on the law of large numbers and the law of iterated logarithm, see [2, 13]. It holds that using n samples this approach gives the searched value with an absolute error of order $O(((\log \log n)/n)^{1/2})$, which practically behaves as $O(n^{-1/2})$. Since in all our simulations we will work with $n = 10^8$, our first approaches to the desired probabilities will have an absolute error of order 10^{-4} .
- **Step 2:** We improve the previous Monte Carlo estimations by using certain linear constraints among the searched probabilities, proved in this work. We take as final estimation of the searched probabilities the least squares solution ([15, Def. 6.1]) of the inconsistent overdetermined system obtained when the Monte Carlo's observed relative frequencies are forced to satisfy these linear constraints. See Section 3.2 for more details.

To have a flavour of the type of results that we will obtain we describe several consequences of some of our results for linear homogeneous differential or difference equations of order n with constant coefficients, see Sections 5 and 7. A first result is that in both cases the expected stability index is $n/2$. Moreover, let r_n denote the probability of the 0 solution to be a global stable attractor (stability index equals n) for them. Then, for differential equations, $r_n \leq 1/2^n$. Furthermore, $r_1 = 1/2$, $r_2 = 1/4$, $r_3 = 1/16$ and our two steps approach gives that $r_4 \simeq 0.00925$, $r_5 \simeq 0.00071$, and that r_k is smaller than 10^{-4} for bigger k . In the case of difference equations we prove that $r_1 = 1/2$ and $r_2 = \frac{1}{\pi} \arctan(\sqrt{2}) \simeq 0.304$.

2 A suitable probability space

In our approach, the starting point is to determine which is the “natural” election of the probability space and the distribution law of the coefficients of the linear dynamical system.

Only after this step is fixed we can ask for the probabilities of some dynamical features or some phase portraits.

For completeness, we start with some previous considerations and with an example, already considered in the literature, see [1, 11, 18]. Consider the planar linear differential system:

$$\begin{pmatrix} \dot{x} \\ \dot{y} \end{pmatrix} = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} \quad (2)$$

where A, B, C, D are random variables, so that the sample space is $\Omega = \mathbb{R}^4$. It is plausible to require that these real random variables are i.i.d. and continuous. Also, according to the *principle of indifference* (or principle of insufficient reason) [6], it would seem reasonable to impose that them were such that the random vector (A, B, C, D) had some kind of uniform distribution in \mathbb{R}^4 . But there is no uniform distribution for unbounded probability spaces. However still, there is a natural election for the distribution of the variables A, B, C and D .

Indeed, it is well known that the phase portrait of the above system does not vary if we multiply the right-hand side of both equations by a positive constant (which corresponds to a constant time scale). This means that in the space of parameters, \mathbb{R}^4 , all the systems with parameters belonging to the same half-straight line passing through the origin are topologically equivalent and in particular have the same stability index. Hence, we can ask for a probability distribution density f of the coefficients such that the random vector

$$\left(\frac{A}{S}, \frac{B}{S}, \frac{C}{S}, \frac{D}{S} \right), \quad \text{with } S = \sqrt{A^2 + B^2 + C^2 + D^2}, \quad (3)$$

has a uniform density on the sphere $\mathbb{S}^3 \subset \mathbb{R}^4$, that is a compact set.

The question is, which are the probability densities f that give rise to a uniform distribution of the vector (3) on the sphere? The answer is that, just assuming that f is continuous and positive, f must be the density of a normal random variable with zero mean. Moreover, this result is true for arbitrary dimension, see next theorem. We remark that the reciprocal result is well known [12, 14].

Theorem 1. *Let X_1, X_2, \dots, X_n be i.i.d. one-dimensional random variables with a continuous positive density function f . The random vector*

$$\left(\frac{X_1}{S}, \frac{X_2}{S}, \dots, \frac{X_n}{S} \right), \quad \text{with } S = \left(\sum_{i=1}^n X_i^2 \right)^{1/2},$$

has a uniform distribution in $\mathbb{S}^{n-1} \subset \mathbb{R}^n$ if and only if each X_i is a normal random variable with zero mean.

Curiously, in the case that we cannot assign uniform distributions, there is an extension of the indifference principle which suggest to use those distributions that maximize the

entropy, i.e. the quantity $h(f) = - \int_{\Omega} f(x) \ln(f(x)) dx$ for any given density f . The one-dimensional random variables with continuous probability density function f on $\Omega = \mathbb{R}$ that maximize the entropy are again the Gaussian ones, [6, Thm 3.2].

Of course, if instead of properties concerning general dynamical systems one focuses on particular models in which the parameters have specific restrictions due to physical or biological reasons one must consider other type of distributions, see for instance [16].

Using Theorem 1, and going back to the initial motivating example, in order to study (2) we have to consider the probability space (Ω, \mathcal{F}, P) where $\Omega = \mathbb{R}^4$, \mathcal{F} is the σ -algebra generated by the open sets of \mathbb{R}^4 and $P : \mathcal{F} \rightarrow [0, 1]$ is the probability function with density $\frac{1}{4\pi^2} e^{-(a^2+b^2+c^2+d^2)/2}$, where for simplicity we take standard deviation 1.

For instance, assume that we want to compute the probability α of system (2) to have exactly one eigenvalue with negative real part. Since the probability of having one null eigenvalue is zero (because events such that in its characterization appears an algebraic equality have zero probability), we have that α coincides with the probability of having a saddle (stability index 1) at the origin, i.e. $AD - BC < 0$. Then, the open set $\mathcal{U} := \{(a, b, c, d) \in \mathbb{R}^4 : ad - bc < 0\}$ belongs to \mathcal{F} and

$$\alpha = P(AD - BC < 0) = \frac{1}{4\pi^2} \int_{\mathcal{U}} e^{-\frac{a^2+b^2+c^2+d^2}{2}} da db dc dd,$$

that is $1/2$ by symmetry, as we will see.

Proof of Theorem 1. Let (X_1, \dots, X_n) be the random vector associated with the with the random variables of the statement, with joint continuous density function $g(x_1, \dots, x_n)$. We claim that

$$g(x_1, \dots, x_n) = h(x_1^2 + \dots + x_n^2), \tag{4}$$

for some continuous function h .

Taking spherical coordinates, we consider the new random vector $(R, \Theta) \in \mathbb{R}^n$ where $R = (X_1^2 + \dots + X_n^2)^{1/2}$ and $\Theta = (\Theta_1, \dots, \Theta_{n-1})$. We have $X_1 = R \cos \Theta_1$, $X_2 = R \sin \Theta_1 \cos \Theta_2, \dots$, $X_{n-1} = R \sin \Theta_1 \sin \Theta_2 \dots \sin \Theta_{n-2} \cos \Theta_{n-1}$ and $X_n = R \sin \Theta_1 \sin \Theta_2 \dots \sin \Theta_{n-2} \sin \Theta_{n-1}$. By the change of variables theorem the joint density function of (R, Θ) is

$$g_{R,\Theta}(r, \theta) = g(r \cos(\theta_1), \dots, r \sin(\theta_1) \dots \sin(\theta_{n-1})) r^{n-1} \sin^{n-2}(\theta_1) \sin^{n-1}(\theta_2) \dots \sin(\theta_{n-2}) \cdot \chi$$

where $\theta = (\theta_1, \dots, \theta_{n-1})$, and

$$\chi := \chi_{[0,\infty)}(r) \cdot \chi_{[0,2\pi)}(\theta_{n-1}) \cdot \prod_{i=1}^{n-2} \chi_{[0,\pi)}(\theta_i),$$

where χ_A stands for the characteristic function of the set A .

The density function of (R, Θ) conditioned to R , $g_{\Theta|R}$, is

$$g_{\Theta|R}(r, \theta) := \frac{g_{R, \Theta}(r, \theta)}{g_R(r)}$$

where $g_R(r)$ is the marginal density of R :

$$g_R(r) := \int_0^\pi \cdots \int_0^\pi \int_0^{2\pi} g(r \cos(\theta_1), \dots, r \sin(\theta_1) \cdots \sin(\theta_{n-1})) d\mathcal{S},$$

being $d\mathcal{S} = r^{n-1} \sin^{n-2}(\theta_1) \sin^{n-1}(\theta_2) \cdots \sin(\theta_{n-2}) d\theta_{n-1} \cdots d\theta_1$ the n -dimensional surface element in spherical coordinates.

To prove the statement, we need to characterize which are the joint density functions $g(x_1, \dots, x_n)$ such that when we fix $R = r$, the probability on the $(n-1)$ -dimensional sphere of radius r , denoted by $\mathbb{S}^{n-1}(r)$, is uniformly distributed. In such a case the partial spherical segment $\Sigma_r = \{R = r, \theta_i \in [\alpha_i, \beta_i] \text{ for } i = 1, \dots, n-1\}$ must have probability $P(\Sigma_r) = \mathcal{S}(\Sigma_r) / \mathcal{S}(\mathbb{S}^{n-1}(r))$ where \mathcal{S} denotes the surface area. Set $\alpha = (\alpha_1, \dots, \alpha_{n-1})$ and $\beta = (\beta_1, \dots, \beta_{n-1})$. Notice that

$$\mathcal{S}(\Sigma_r) = \int_{\alpha_1}^{\beta_1} \cdots \int_{\alpha_{n-1}}^{\beta_{n-1}} d\mathcal{S} = r^{n-1} A(\alpha, \beta)$$

where

$$A(\alpha, \beta) = \int_{\alpha_1}^{\beta_1} \cdots \int_{\alpha_{n-1}}^{\beta_{n-1}} \sin^{n-2}(\theta_1) \sin^{n-1}(\theta_2) \cdots \sin(\theta_{n-2}) d\theta_{n-1} \cdots d\theta_1$$

and $\mathcal{S}(\mathbb{S}^{n-1}(r)) = \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})} r^{n-1}$. Hence, on the one hand,

$$P(\Sigma_r) = \Gamma\left(\frac{n}{2}\right) \frac{A(\alpha, \beta)}{2\pi^{\frac{n}{2}}},$$

which does not depend on r . On the other hand,

$$P(\Sigma_r) = \int_{\alpha_1}^{\beta_1} \cdots \int_{\alpha_{n-1}}^{\beta_{n-1}} g_{\Theta|R} d\theta$$

where $d\theta = d\theta_{n-1} \cdots d\theta_2 d\theta_1$. This implies that

$$\begin{aligned} & \int_{\alpha_1}^{\beta_1} \cdots \int_{\alpha_{n-1}}^{\beta_{n-1}} \frac{g(r \cos(\theta_1), \dots, r \sin(\theta_1) \cdots \sin(\theta_{n-1})) r^{n-1} \sin^{n-2}(\theta_1) \sin^{n-1}(\theta_2) \cdots \sin(\theta_{n-2}) \cdot \chi}{g_R(r)} d\theta \\ &= \frac{\Gamma\left(\frac{n}{2}\right)}{2\pi^{\frac{n}{2}}} \int_{\alpha_1}^{\beta_1} \cdots \int_{\alpha_{n-1}}^{\beta_{n-1}} \sin^{n-2}(\theta_1) \sin^{n-1}(\theta_2) \cdots \sin(\theta_{n-2}) d\theta, \end{aligned}$$

for all $\alpha_i, \beta_i \in [0, \pi)$ for $i = 1, \dots, n-2$ with $\alpha_i < \beta_i$ and $\alpha_{n-2}, \beta_{n-2} \in [0, 2\pi)$ with $\alpha_{n-2} < \beta_{n-2}$. This last equality implies that almost everywhere

$$\frac{\Gamma\left(\frac{n}{2}\right)}{2\pi^{\frac{n}{2}}} = \frac{g(r \cos(\theta_1), \dots, r \sin(\theta_1) \cdots \sin(\theta_{n-1})) r^{n-1}}{g_R(r)}$$

and therefore $g(r \cos(\theta_1), \dots, r \sin(\theta_1) \cdots \sin(\theta_{n-1}))$ is a function that only depends on r . In consequence, writing this fact in cartesian coordinates, we get that almost everywhere $g(x_1, \dots, x_n) = h(x_1^2 + \cdots + x_n^2)$, for some continuous function h and the claim (4) follows.

Now we complete the proof. Since X_1, \dots, X_n are i.i.d., with positive density f , we know that $g(x_1, \dots, x_n) = f(x_1) \dots f(x_n)$. So equation (4) writes as

$$f(x_1) \cdots f(x_n) = h(x_1^2 + \cdots + x_n^2) \quad \text{for all } (x_1, \dots, x_n) \in \mathbb{R}^n$$

and $h > 0$. Taking $x_2 = \cdots = x_n = 0$ we have that $f(x_1) f(0)^{n-1} = h(x_1^2)$ and $h(0) = (f(0))^n > 0$. Thus,

$$f(x_1) \cdots f(x_n) = \frac{h(x_1^2)}{(f(0))^{n-1}} \cdots \frac{h(x_n^2)}{(f(0))^{n-1}} = \frac{h(x_1^2)}{(f(0))^n} \cdots \frac{h(x_n^2)}{(f(0))^n} (f(0))^n = h(x_1^2 + \cdots + x_n^2).$$

Hence, using that $h(0) = (f(0))^n > 0$,

$$\frac{h(x_1^2)}{h(0)} \cdots \frac{h(x_n^2)}{h(0)} = \frac{h(x_1^2 + \cdots + x_n^2)}{h(0)}.$$

Taking $H(\xi) := h(\xi)/h(0)$, and $u_i = x_i^2$, it holds that

$$H(u_1) \cdots H(u_n) = H(u_1 + \cdots + u_n) \quad \text{with } H(0) = 1. \quad (5)$$

Hence, $\varphi(U) = \log(H(u))$ is a continuous function that satisfies the Cauchy's functional equation

$$\varphi(u_1) + \cdots + \varphi(u_n) = \varphi(u_1 + \cdots + u_n) \quad \text{with } \varphi(0) = 0.$$

It is well known that all its continuous solutions are $\varphi(x) = ax$, for some $a \in \mathbb{R}$. Hence all continuous solutions of (5) are $H(x) = e^{ax}$.

As a consequence, $f(x) = b e^{ax^2}$ for some $(a, b) \in \mathbb{R}^2$. Since f is a density function, $a < 0$. Moreover, using $\int_{-\infty}^{\infty} b e^{ax^2} dx = b \sqrt{-\pi/a} = 1$, and setting $a = -1/(2\sigma^2)$, we get that

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{x^2}{2\sigma^2}},$$

so each variable X_i is a normal random variable $N(0, \sigma^2)$.

The reciprocal part is straightforward and well known [12, 14]. ■

Remark 1. The continuity condition for f in Theorem 1 is relevant since Equation (5) also admits non-continuous solutions that can be constructed, for instance, from non-continuous solutions of the Cauchy's functional equation known for $n = 2$, see [10].

3 A preliminary result and methodology

We will investigate the probabilities of having a certain stability index for several linear dynamical systems with random coefficients. In particular we consider:

- (a) Differential systems $\dot{\mathbf{x}} = A \mathbf{x}$ where $\mathbf{x} \in \mathbb{R}^n$ and A is a real constant $n \times n$ matrix,
- (b) Homogeneous linear differential equation of order n with constant coefficients: $a_n x^{(n)} + a_{n-1} x^{(n-1)} + \dots + a_1 x' + a_0 x = 0$,
- (c) Linear discrete systems $b \mathbf{x}_{k+1} = A \mathbf{x}_k$ where $\mathbf{x}_k \in \mathbb{R}^n$, $b \in \mathbb{R}$; and A is a real constant $n \times n$ matrix,
- (d) Linear homogeneous difference equation of order n with constant coefficients $a_n x_{k+n} + a_{n-1} x_{k+n-1} + \dots + a_1 x_{k+1} + a_0 x_k = 0$.

Notice that in the four situations the behaviour of the dynamical systems does not change if we multiply all the involved constants by the same positive real number. This fact situates the four problems in the same context that the motivating example (2). Hence, following the results of Section 2, in all the cases the coefficients will be i.i.d. normal variables with zero mean and standard deviation 1.

Hence in all cases we have a well-defined probability space (Ω, \mathcal{F}, P) , where $\Omega = \mathbb{R}^m$, with $m = n^2, n+1, n^2+1$ or $n+1$ according we are in case (a), (b), (c) or (d), respectively, \mathcal{F} is the σ -algebra generated by the open sets and for each $\mathcal{A} \in \mathcal{F}$,

$$P(\mathcal{A}) = \frac{1}{(\sqrt{2\pi})^m} \int_{\mathcal{A}} e^{-\|\mathbf{a}\|^2/2} d\mathbf{a},$$

where $\mathbf{a} = (a_1, a_2, \dots, a_m)$, $\|\mathbf{a}\|^2 = \sum_{j=1}^m a_j^2$ and $d\mathbf{a} = da_1 da_2 \dots da_m$. For instance the matrices A appearing in case (a) and (c) are the so called *random matrices*.

The use of Routh-Hurwitz algorithm is a very useful tool to count the number of roots of a polynomial with negative real parts and it is implemented in many computer algebra systems. These conditions are given in terms of algebraic inequalities among the coefficients of the polynomials. Let us recall how to use it to count the number of roots with modulus less than one of a polynomial and, hence, to obtain the so called Jury conditions.

Given any polynomial $Q(\lambda) = q_n \lambda^n + q_{n-1} \lambda^{n-1} + \dots + q_1 \lambda + q_0$ with $q_j \in \mathbb{C}$, by using the conformal transformation $\lambda = \frac{z+1}{z-1}$, we get the associated polynomial

$$Q^*(z) = q_n (z+1)^n + q_{n-1} (z+1)^{n-1} (z-1) + \dots + q_0 (z-1)^n. \quad (6)$$

It is straightforward to observe that $\lambda_0 \in \mathbb{C}$ is a root of $Q(\lambda)$ such that $|\lambda_0| < 1$ if and only if $z_0 = (\lambda_0 + 1)/(\lambda_0 - 1)$ is a root of $Q^*(z)$ such that $\text{Re}(z_0) < 0$.

Hence, because Routh-Hurwitz and Jury conditions are semi-algebraic, in all cases the random variable that X that assigns to each dynamical system its stability index k , $0 \leq k \leq n$, is measurable. Hence $\mathcal{A}_k := \{\mathbf{a} \in \mathbb{R}^m : X(\mathbf{a}) = k\} \in \mathcal{F}$ and its probability $p_k := P(\mathcal{A}_k)$ is well defined. Observe also that the non-hyperbolic cases are totally negligible because for their characterization some algebraic equalities appear. In this paper we will either calculate or estimate in the four situations the values p_k for $k \leq 10$.

3.1 A preliminary result

In three of the above considered cases we will apply the following auxiliary result:

Lemma 2. *Let (Ω, \mathcal{F}, P) be a probability space and let $Y : \Omega \rightarrow \mathbb{R}$ be a discrete random variable with image $\text{Im}(Y) = \{0, 1, \dots, n\}$, and probability function $p_k = P(Y = k)$ such that $p_k = p_{n-k}$ for all $k = 0, \dots, n$. Then $E(Y) = \sum_{k=0}^n kp_k = n/2$. Moreover*

(a) *If n is odd then $2 \sum_{k=0}^{\frac{n-1}{2}} p_k = 1$. In particular, when $n = 1$, $p_0 = p_1 = \frac{1}{2}$.*

(b) *If n is even and $n \geq 2$ then $2 \sum_{k=0}^{\frac{n}{2}-1} p_k + p_{\frac{n}{2}} = 1$.*

If, additionally, n is even and $\sum_{i \text{ odd}} p_i = \sum_{i \text{ even}} p_i = \frac{1}{2}$, then*

(c) *If $\frac{n}{2}$ is even, then $2 \sum_{k=0, k \text{ even}}^{\frac{n}{2}-2} p_k + p_{\frac{n}{2}} = \frac{1}{2}$, and $2 \sum_{k=1, k \text{ odd}}^{\frac{n}{2}-1} p_k = \frac{1}{2}$. In particular, when $n = 4$, $p_1 = p_3 = \frac{1}{4}$, $p_2 = \frac{1}{2} - 2p_0$ and $p_4 = p_0$.*

(d) *If $\frac{n}{2}$ is odd, then $2 \sum_{k=0, k \text{ even}}^{\frac{n}{2}-1} p_k = \frac{1}{2}$, and $\sum_{k=1, k \text{ odd}}^{\frac{n}{2}-2} p_k + p_{\frac{n}{2}} = \frac{1}{2}$. In particular, when $n = 2$, $p_0 = p_2 = \frac{1}{4}$ and $p_1 = \frac{1}{2}$.*

Proof. We start proving that $E(Y) = n/2$. Assume for instance that n is odd. Since $p_k = p_{n-k}$, it holds that $kp_k + (n-k)p_{n-k} = np_k$, for each $k \leq (n-1)/2$. Hence,

$$\begin{aligned} E(X) &= np_0 + np_1 + \dots + np_{\frac{n-1}{2}} = \frac{n}{2} \left(2p_0 + 2p_1 + \dots + 2p_{\frac{n-1}{2}} \right) \\ &= \frac{n}{2} \left((p_0 + p_n) + (p_1 + p_{n-1}) + \dots + (p_{\frac{n-1}{2}} + p_{\frac{n+1}{2}}) \right) = \frac{n}{2}. \end{aligned}$$

When n is even the proof is similar.

The proof of all the four items is straightforward and we omit it. ■

3.2 Experimental methodology

In all the cases considered in the paper, when we can not give an exact value of the probabilities p_k we start estimating them by using the *Monte Carlo* method (further details

*When n is odd the imposed equalities automatically hold.

are given in each of the following subsections), [13]. From these obtained estimations, the observed relative frequencies, we improve them via the *least squares* method, by using the linear constraints given in Corollaries 4, 6 and 13.

A more detailed explanation of this second step is as follows: The probabilities p_k satisfy some affine relations, like the ones in Lemma 2 or the ones in Proposition 12. Then, if we denote $\mathbf{p} = (p_0, \dots, p_n)^t \in \mathbb{R}^{n+1}$ it is possible to write $\mathbf{p} = M\mathbf{q} + b$ where $\mathbf{q} \in \mathbb{R}^k$ with $k \leq n$ is a vector whose components are different elements of p_0, \dots, p_n ; $M \in \mathcal{M}_{n \times k}(\mathbb{R})$, and $b \in \mathbb{R}^k$. Let $\tilde{\mathbf{p}} = (\tilde{p}_0, \dots, \tilde{p}_n)^t$ be the vector with the estimated probabilities obtained by the observed relative frequencies using the Monte Carlo method. Then, we can find the least squares solution [15, Def. 6.1] of the the system,

$$\tilde{\mathbf{p}} = M\hat{\mathbf{q}} + b, \quad (7)$$

obtaining that

$$\hat{\mathbf{q}} = (M^t \cdot M)^{-1} \cdot M^t \cdot (\tilde{\mathbf{p}} - b), \quad (8)$$

see [7, p. 198] or [17, p. 200]. So we can find some improved estimations $\hat{\mathbf{p}}$, via:

$$\hat{\mathbf{p}} = M\hat{\mathbf{q}} + b. \quad (9)$$

Some detailed examples are given in Sections 4, 5 and 7.

4 Linear random differential systems

Consider linear differential systems $\dot{\mathbf{x}} = A\mathbf{x}$ where $\mathbf{x} \in \mathbb{R}^n$, $A \in \mathcal{M}_{n \times n}(\mathbb{R})$, where A is a random matrix which entries are i.i.d. random variables with $N(0, 1)$ distribution. Let X be the random variable that counts the number of eigenvalues of A with negative real part, $s(A)$.

Proposition 3. *With the above notations, set $p_k = P(X = k)$. The following holds:*

- (a) $\sum_{k=0}^n p_k = 1$.
- (b) For all $k \in \{0, 1, \dots, n\}$, $p_k = p_{n-k}$.
- (c) $\sum_{i \text{ odd}} p_i = \sum_{i \text{ even}} p_i = \frac{1}{2}$.

Proof. The assertion (a) is trivial. To prove (b) we observe that if a matrix A has k eigenvalues with negative real part, then $B = -A$ has $n - k$ eigenvalues with negative real part. Calling q_m the probability that B has m eigenvalues with negative real part, we get that $p_m = q_m$. This is so, because if $X \sim N(0, 1)$ then $-X \sim N(0, 1)$ and as a consequence the entries of A and B have the same distribution. Then, $q_k = p_{n-k}$ and the result follows.

To see (c) we claim that $s(A)$ is even if and only if the determinant of A is positive and, moreover, $P(\det(A) > 0) = 1/2$. From this claim we get the result because $\sum_{i \text{ even}} p_i$ is the probability of $s(A)$ to be even. To prove the first part of the claim notice first that we can assume that $0 \neq \det(A) = \lambda_1 \lambda_2 \cdots \lambda_n$, where $\lambda_1, \lambda_2, \dots, \lambda_n$ are the n eigenvalues of A . We write $\lambda_1 \lambda_2 \cdots \lambda_n = (\lambda_1 \lambda_2 \cdots \lambda_k)(\lambda_{k+1} \lambda_{k+2} \cdots \lambda_n)$ where $\lambda_1, \lambda_2, \dots, \lambda_k$ are all the real negative ones. Observe also that for complex eigenvalues $\lambda \bar{\lambda} > 0$. Hence $\lambda_{k+1} \lambda_{k+2} \cdots \lambda_n > 0$, $\text{sign}(\det(A)) = (-1)^k$ and the condition $s(A)$ even is characterized by $\det(A) > 0$. To prove that $P(\det(A) > 0) = 1/2$ note that if B is the matrix obtained by changing the sign of one column of A then $\det(A) \cdot \det(B) < 0$ and hence $P(\det(A) < 0) = P(\det(B) > 0)$. Furthermore, since the entries of A and B have the same distribution we have $P(\det(B) > 0) = P(\det(A) > 0)$, thus $P(\det(A) < 0) = P(\det(A) > 0) = 1/2$. ■

From the above proposition it easily follows:

Corollary 4. *Consider $\dot{\mathbf{x}} = A\mathbf{x}$, $\mathbf{x} \in \mathbb{R}^n$ with $A \in \mathcal{M}_{n \times n}(\mathbb{R})$ a random matrix with i.i.d. $N(0, 1)$ entries, let X be the random variable defined above and $p_k = P(X = k)$. Then the probabilities p_k satisfy all the consequences of Lemma 2. In particular $E(X) = n/2$.*

Now we reproduce some experiments to estimate the probabilities p_k for low dimensional cases. We apply the Monte Carlo method, that is, for each considered dimension n , we have generated 10^8 matrices $A \in \mathcal{M}_{n \times n}(\mathbb{R})$ whose entries are pseudorandom numbers simulating the realizations on n^2 independent random variables with $N(0, 1)$ distribution. For each matrix A we have computed the characteristic polynomial, and counted the number of eigenvalues with negative real part by using the Routh-Hurwitz zeros counter [8, p. 1076]. For $n \geq 5$ and in order to decrease the computation time we have directly computed numerically the eigenvalues of A and counted the number of them with negative real part.

For each considered dimension of the phase space n , and in order to take advantage of the relations stated in Corollary 4, we can refine the solutions using the least squares solutions of the inconsistent linear system associated to these relations when using the observed frequencies obtained by the Monte Carlo simulation.

We detail one example. Set $n = 7$, for instance. By Corollary 4 we have $p_3 = p_4 = \frac{1}{2} - p_0 - p_1 - p_2$; $p_5 = p_2$; $p_6 = p_1$ and $p_7 = p_0$. So, using the notation introduced in

Section 3.2, we can write $\mathbf{p} = M\mathbf{q} + b$, where $\mathbf{p}^t = (p_0, \dots, p_7)$;

$$M = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ -1 & -1 & -1 \\ -1 & -1 & -1 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}; \quad \mathbf{q} = \begin{pmatrix} p_0 \\ p_1 \\ p_2 \end{pmatrix}; \quad \text{and } b = \begin{pmatrix} 0 \\ 0 \\ 0 \\ \frac{1}{2} \\ \frac{1}{2} \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

The observed relative frequencies in our Monte Carlo simulation are

$$\tilde{\mathbf{p}}^t = \left(\frac{31643}{50000000}, \frac{261137}{12500000}, \frac{7124967}{50000000}, \frac{1344047}{4000000}, \frac{33597117}{100000000}, \frac{14248187}{100000000}, \frac{1043913}{50000000}, \frac{63379}{100000000} \right).$$

By finding the least squares solution of the system (7) ([7, p. 198] or [17, p. 200]), given by (9), we obtain

$$\hat{\mathbf{p}}^t = \left(\frac{25333}{40000000}, \frac{2088461}{100000000}, \frac{28498121}{200000000}, \frac{16799573}{50000000}, \frac{16799573}{50000000}, \frac{28498121}{200000000}, \frac{2088461}{100000000}, \frac{25333}{40000000} \right).$$

The other cases follow similarly.

We summarize the results of our experiments in the Table 1, where the observed relative frequencies and the estimates are presented only up to the fifth decimal (in the table, and in the whole paper, frequency stands for relative frequency) because as we already explained in the introduction, the expected absolute error will be of order 10^{-4} .

Dimension	Observed frequency	Least squares	Relations (Corol. 4)
$n = 1$	$\tilde{p}_0 = 0.49996$ $\tilde{p}_1 = 0.50004$		$p_0 = 0.5$ $p_1 = 0.5$
$n = 2$	$\tilde{p}_0 = 0.24999$ $\tilde{p}_1 = 0.50006$ $\tilde{p}_2 = 0.24995$		$p_0 = 0.25$ $p_1 = 0.5$ $p_2 = 0.25$
$n = 3$	$\tilde{p}_0 = 0.10447$ $\tilde{p}_1 = 0.39542$ $\tilde{p}_2 = 0.39557$ $\tilde{p}_3 = 0.10454$	$\hat{p}_0 = 0.10450$ $\hat{p}_1 = 0.39550$ $\hat{p}_2 = 0.39550$ $\hat{p}_3 = 0.10450$	p_0 $p_1 = \frac{1}{2} - p_0$ $p_2 = \frac{1}{2} - p_0$ $p_3 = p_0$
$n = 4$	$\tilde{p}_0 = 0.03722$ $\tilde{p}_1 = 0.25009$ $\tilde{p}_2 = 0.42556$ $\tilde{p}_3 = 0.24998$ $\tilde{p}_4 = 0.03715$	$\hat{p}_0 = 0.03721$ $\hat{p}_1 = 0.25000$ $\hat{p}_2 = 0.42558$ $\hat{p}_3 = 0.25000$ $\hat{p}_4 = 0.03721$	p_0 $p_1 = \frac{1}{4}$ $p_2 = \frac{1}{2} - 2p_0$ $p_3 = \frac{1}{4}$ $p_4 = p_0$
$n = 5$	$\tilde{p}_0 = 0.01126$ $\tilde{p}_1 = 0.13028$ $\tilde{p}_2 = 0.35848$ $\tilde{p}_3 = 0.35852$ $\tilde{p}_4 = 0.13020$ $\tilde{p}_5 = 0.01126$	$\hat{p}_0 = 0.01126$ $\hat{p}_1 = 0.13024$ $\hat{p}_2 = 0.35850$ $\hat{p}_3 = 0.35850$ $\hat{p}_4 = 0.13024$ $\hat{p}_5 = 0.01126$	p_0 p_1 $p_2 = \frac{1}{2} - p_0 - p_1$ $p_3 = \frac{1}{2} - p_0 - p_1$ $p_4 = p_1$ $p_5 = p_0$
$n = 6$	$\tilde{p}_0 = 0.00289$ $\tilde{p}_1 = 0.05675$ $\tilde{p}_2 = 0.24710$ $\tilde{p}_3 = 0.38642$ $\tilde{p}_4 = 0.24714$ $\tilde{p}_5 = 0.56810$ $\tilde{p}_6 = 0.00289$	$\hat{p}_0 = 0.00288$ $\hat{p}_1 = 0.05678$ $\hat{p}_2 = 0.24712$ $\hat{p}_3 = 0.38644$ $\hat{p}_4 = 0.24712$ $\hat{p}_5 = 0.05678$ $\hat{p}_6 = 0.00288$	p_0 p_1 $p_2 = \frac{1}{4} - p_0$ $p_3 = \frac{1}{2} - 2p_1$ $p_4 = \frac{1}{4} - p_0$ $p_5 = p_1$ $p_6 = p_0$
$n = 7$	$\tilde{p}_0 = 0.00063$ $\tilde{p}_1 = 0.02089$ $\tilde{p}_2 = 0.14250$ $\tilde{p}_3 = 0.33601$ $\tilde{p}_4 = 0.33597$ $\tilde{p}_5 = 0.14248$ $\tilde{p}_6 = 0.02088$ $\tilde{p}_7 = 0.00063$	$\hat{p}_0 = 0.00063$ $\hat{p}_1 = 0.02088$ $\hat{p}_2 = 0.14249$ $\hat{p}_3 = 0.33600$ $\hat{p}_4 = 0.33600$ $\hat{p}_5 = 0.14249$ $\hat{p}_6 = 0.02088$ $\hat{p}_7 = 0.00063$	p_0 p_1 p_2 $p_3 = \frac{1}{2} - p_0 - p_1 - p_2$ $p_4 = \frac{1}{2} - p_0 - p_1 - p_2$ $p_5 = p_2$ $p_6 = p_1$ $p_7 = p_0$

Dimension	Observed frequency	Least squares	Relations (Corol. 4)
$n = 8$	$\tilde{p}_0 = 0.00012$ $\tilde{p}_1 = 0.00651$ $\tilde{p}_2 = 0.06948$ $\tilde{p}_3 = 0.24356$ $\tilde{p}_4 = 0.36080$ $\tilde{p}_5 = 0.24346$ $\tilde{p}_6 = 0.06946$ $\tilde{p}_7 = 0.00650$ $\tilde{p}_8 = 0.00012$	$\hat{p}_0 = 0.00012$ $\hat{p}_1 = 0.00650$ $\hat{p}_2 = 0.06948$ $\hat{p}_3 = 0.24350$ $\hat{p}_4 = 0.36080$ $\hat{p}_5 = 0.24350$ $\hat{p}_6 = 0.06948$ $\hat{p}_7 = 0.00650$ $\hat{p}_8 = 0.00012$	p_0 p_1 p_2 $p_3 = \frac{1}{4} - p_1$ $p_4 = \frac{1}{2} - 2p_0 - 2p_2$ $p_5 = \frac{1}{4} - p_1$ $p_6 = p_2$ $p_7 = p_1$ $p_8 = p_0$
$n = 9$	$\tilde{p}_0 = 0.00002$ $\tilde{p}_1 = 0.00171$ $\tilde{p}_2 = 0.02880$ $\tilde{p}_3 = 0.14952$ $\tilde{p}_4 = 0.31987$ $\tilde{p}_5 = 0.31999$ $\tilde{p}_6 = 0.14958$ $\tilde{p}_7 = 0.02878$ $\tilde{p}_8 = 0.00171$ $\tilde{p}_9 = 0.00002$	$\hat{p}_0 = 0.00002$ $\hat{p}_1 = 0.00171$ $\hat{p}_2 = 0.02879$ $\hat{p}_3 = 0.14955$ $\hat{p}_4 = 0.31993$ $\hat{p}_5 = 0.31993$ $\hat{p}_6 = 0.14955$ $\hat{p}_7 = 0.02879$ $\hat{p}_8 = 0.00171$ $\hat{p}_9 = 0.00002$	p_0 p_1 p_2 p_3 $p_4 = \frac{1}{2} - p_0 - p_1 - p_2 - p_3$ $p_5 = \frac{1}{2} - p_0 - p_1 - p_2 - p_3$ $p_6 = p_3$ $p_7 = p_2$ $p_8 = p_1$ $p_9 = p_0$
$n = 10$	$\tilde{p}_0 = 0.00000$ $\tilde{p}_1 = 0.00038$ $\tilde{p}_2 = 0.01015$ $\tilde{p}_3 = 0.07850$ $\tilde{p}_4 = 0.23987$ $\tilde{p}_5 = 0.34224$ $\tilde{p}_6 = 0.23984$ $\tilde{p}_7 = 0.07849$ $\tilde{p}_8 = 0.01015$ $\tilde{p}_9 = 0.00038$ $\tilde{p}_{10} = 0.00000$	$\hat{p}_0 = 0.00000$ $\hat{p}_1 = 0.00038$ $\hat{p}_2 = 0.01015$ $\hat{p}_3 = 0.07850$ $\hat{p}_4 = 0.23985$ $\hat{p}_5 = 0.34224$ $\hat{p}_6 = 0.23985$ $\hat{p}_7 = 0.07850$ $\hat{p}_8 = 0.01015$ $\hat{p}_9 = 0.00038$ $\hat{p}_{10} = 0.00000$	p_0 p_1 p_2 p_3 $p_4 = \frac{1}{4} - p_0 - p_2$ $p_5 = \frac{1}{2} - 2p_1 - 2p_3$ $p_6 = \frac{1}{4} - p_0 - p_2$ $p_7 = p_3$ $p_8 = p_2$ $p_9 = p_1$ $p_{10} = p_0$

Table 1. Linear stability indices for linear random differential systems.

5 Linear random differential equations of order n

In this section we consider linear random homogeneous differential equations of order n

$$A_n x^{(n)} + A_{n-1} x^{(n-1)} + \cdots + A_2 x'' + A_1 x' + A_0 x = 0, \quad (10)$$

where $x = x(t)$, the derivatives are taken in respect to t , and A_j are again i.i.d. random variables with $N(0, 1)$ distribution.

To get the stability index for them we only need to know the probability distributions of the number of roots with negative real part of its associated random characteristic polynomial:

$$Q(\lambda) = A_n \lambda^n + A_{n-1} \lambda^{n-1} + \cdots + A_1 \lambda + A_0.$$

Let X be the random variable that counts the number of roots of $Q(\lambda)$ with negative real parts and define $p_k = P(X = k)$ for $k = 0, 1, \dots, n$.

Proposition 5. *Set $p_k = P(X = k)$, where X is the random variable defined above. Then*

- (a) $\sum_{k=0}^n p_k = 1$.
- (b) For all $k \in \{0, 1, \dots, n\}$, $p_k = p_{n-k}$.
- (c) $\sum_{i \text{ odd}} p_i = \sum_{i \text{ even}} p_i = \frac{1}{2}$.

Proof. The proof of (a) is trivial. To prove (b) consider equation (10) with its characteristic polynomial $Q(\lambda)$ and also the new differential equation

$$(-1)^n A_n x^{(n)} + (-1)^{n-1} A_{n-1} x^{(n-1)} + \cdots + A_2 x'' - A_1 x' + A_0 x = 0 \quad (11)$$

with its characteristic polynomial $Q^*(\lambda) = Q(-\lambda) = (-1)^n A_n \lambda^n + (-1)^{n-1} A_{n-1} \lambda^{n-1} + \cdots - A_1 \lambda + A_0$. Since $Q(\lambda) = 0$ if and only if $Q^*(-\lambda) = 0$ we get that $p_k = p_{n-k}^*$ where p_i^* the probability that $Q^*(\lambda)$ has i roots with negative real part. But also $p_k = p_k^*$ because the coefficients of the equations (10) and (11) have the same distributions. Hence, the result follows.

Similarly as in the proof of (c) of Proposition 3 we observe that the polynomial $Q(\lambda)$ has an odd number of roots with negative real part if and only if $A_0 \cdot A_n < 0$, because we can neglect the case of having some roots with zero real part. Since the coefficients of (10) are symmetric independent random variables, the probability that $Q(\lambda)$ has an odd number of roots with negative real part is

$$P(\{A_0 > 0\} \cap \{A_n < 0\}) + P(\{A_0 < 0\} \cap \{A_n > 0\}) = \frac{1}{2} \times \frac{1}{2} + \frac{1}{2} \times \frac{1}{2} = \frac{1}{2}.$$

■

Corollary 6. *Consider the linear random homogeneous differential equation of order n (10), with all A_i being i.i.d. $N(0,1)$ random variables, let X be defined above, and set $p_k = P(X = k)$. Then the probabilities p_k satisfy all the consequences of Lemma 2. In particular $E(X) = n/2$.*

For each n , let r_n be the probability of the origin to be a global stable attractor (asymptotically stable equilibrium) for (10), that is $r_n = p_n$. By Proposition 5(b) this probability coincides with the one of being a repeller because $p_n = p_0$. Our results in Proposition 8 seem to indicate that r_n decreases with n . Before proving this proposition we need a preliminary result.

Lemma 7. *Let U, V, S and T be i.i.d. random variables with normal standard distribution. Then $p^+ := P(U > 0; V > 0; S > 0; T > 0; UT - SV > 0) = 1/32$.*

Proof. Set $\mathcal{A}^\pm = \{U > 0; V > 0; S > 0; T > 0; \pm(UT - SV) > 0\}$, and $\mathcal{A}^0 = \{U > 0; V > 0; S > 0; T > 0; UT - SV = 0\}$. Denote by $p^\pm = P(\mathcal{A}^\pm)$ and $p^0 = P(\mathcal{A}^0)$. Then, since $p^0 = 0$ and $\mathcal{A}^- \cup \mathcal{A}^0 \cup \mathcal{A}^+ = \{U > 0; V > 0; S > 0; T > 0\}$ it holds that $p^+ + p^- = (1/2)^4 = 1/16$. To end the proof it suffices to show that $p^+ = p^-$.

Notice first that

$$\begin{aligned} \mathcal{A}^+ &= \{U > 0; V > 0; S > 0; T > 0; UT - SV > 0\} = \{V > 0; S > 0; T > 0; UT - SV > 0\}, \\ \mathcal{A}^- &= \{U > 0; V > 0; S > 0; T > 0; UT - SV < 0\} = \{U > 0; S > 0; T > 0; SV - UT > 0\}. \end{aligned}$$

This is so, because for instance in the definition of \mathcal{A}^+ , the last inequality can also be written as $U > SV/T > 0$ and from it we know that the condition $U > 0$ can be removed. Finally, interchanging U and V and S and T we get the same relations in the definitions of \mathcal{A}^+ and \mathcal{A}^- . Since all variables are independent $N(0,1)$, both sets have the same probability and $p^+ = p^-$, as we wanted to prove. ■

Proposition 8. *With the above notations, $r_n \leq 1/2^n$, for all $n \geq 1$. Moreover, $r_1 = 1/2$, $r_2 = 1/4$, $r_3 = 1/16$ and $r_4 < r_3/2 = 1/32$.*

Proof of Proposition 8. Notice that r_n is the probability that the characteristic polynomial $Q(\lambda)$, associated to the random differential equation (10) is a Hurwitz stable polynomial, that is $r_n = P(\text{Every root of } Q(\lambda) \text{ belongs to } \mathfrak{R}^-)$, where $\mathfrak{R}^- = \{z \in \mathbb{C} \text{ such that } \text{Re}(z) < 0\}$. It is well-known that a necessary condition for a polynomial to have every root in \mathfrak{R}^- is that all its coefficients have the same sign. This is so because it holds for polynomials of degree 1 and 2, and this property is preserved when we multiply two polynomials satisfying

it. Hence,

$$\{A_0, \dots, A_n \text{ such that all roots of } P(\lambda) \text{ are in } \mathfrak{R}^-\} \subset \left\{ \bigcap_{i=0}^n \{A_i < 0\} \right\} \cup \left\{ \bigcap_{i=0}^n \{A_i > 0\} \right\}. \quad (12)$$

Since the variables A_i are independent and symmetric

$$P\left(\bigcap_{i=0}^n \{A_i < 0\}\right) = P\left(\bigcap_{i=0}^n \{A_i > 0\}\right) = \frac{1}{2^{n+1}}.$$

As a consequence,

$$r_n \leq P\left(\bigcap_{i=0}^n \{A_i < 0\}\right) + P\left(\bigcap_{i=0}^n \{A_i > 0\}\right) = \frac{1}{2^n},$$

and the first statement follows.

The equalities $r_1 = 1/2$ and $r_2 = 1/4$ are a simple consequence that for $n = 1, 2$ the inclusion (12) is an equality.

Let us prove that $r_3 = p_3 = 1/16$. By using the Routh-Hurwitz criterion [8, p. 1076], it can be seen that $a_3\lambda^3 + a_2\lambda^2 + a_1\lambda + a_0$ has every root in \mathfrak{R}^- if and only if all its coefficients have the same sign and moreover $a_1a_2 - a_0a_3 > 0$. Hence, $p_3 = p_3^- + p_3^+$, where $p_3^- := P(A_0 < 0; A_1 < 0; A_2 < 0; A_3 < 0; A_1A_2 - A_0A_3 > 0)$; and $p_3^+ := P(A_0 > 0; A_1 > 0; A_2 > 0; A_3 > 0; A_1A_2 - A_0A_3 > 0)$, being all $A_i \sim N(0, 1)$ and independent. Due to their symmetry, the random variables A_i and $-A_i$, for $i = 0, \dots, 3$ have the same distribution and hence $p_3^+ = p_3^-$. Therefore $p_3 = 2p_3^+$. The result follows now by Lemma 7, which gives $p_3^+ = 1/32$.

Let us prove that $r_4 < r_3/2$. To compare both probabilities, here it will be more convenient to write the coefficients of the polynomials with subscripts with increasing ordering, that is $q_n(x) = a_0x^n + a_1x^{n-1} + \dots + a_{n-1}x + a_n$. With this notation, which also respects the traditional notation when writing the Hurwitz matrices, and when $a_0 > 0$, the Routh-Hurwitz conditions to have stability index n for $n = 3, 4$ are precisely that the principal minors of next matrices

$$\begin{pmatrix} a_1 & a_3 & 0 \\ a_0 & a_2 & 0 \\ 0 & a_1 & a_3 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} a_1 & a_3 & 0 & 0 \\ a_0 & a_2 & a_4 & 0 \\ 0 & a_1 & a_3 & 0 \\ 0 & a_0 & a_2 & a_4 \end{pmatrix},$$

are positive, where the left-hand one corresponds to $n = 3$ and the other one to $n = 4$. Hence, these conditions when $a_0 > 0$ and for $n = 3$ are: $a_1 > 0, a_1a_2 - a_0a_3 > 0$ and $a_3 > 0$. Similarly, for $n = 4$ are $a_1 > 0, a_1a_2 - a_0a_3 > 0, a_3(a_1a_2 - a_0a_3) - a_4a_1^2 > 0$ and $a_4 > 0$.

Consider now, for $n = 3, 4$, the random polynomials $Q_n(x) = \tilde{A}_0 x^n + \tilde{A}_1 x^{n-1} + \dots + \tilde{A}_{n-1} x + \tilde{A}_n$, where $\tilde{A}_i \sim N(0, 1)$ and are independent (notice that with this notation each coefficient \tilde{A}_i is the coefficient A_{n-i} of the characteristic polynomial). For simplicity we denote with the same name the coefficients of Q_3 and Q_4 although they are different random variables. As above, $r_3 = 2p_3^+$, and $r_4 = 2p_4^+$, where $p_k^+ = P(\mathcal{A}_k^+)$, with

$$\begin{aligned}\mathcal{A}_3^+ &= \{\tilde{A}_0 > 0; \tilde{A}_1 > 0; \tilde{A}_3 > 0; \tilde{A}_1 \tilde{A}_2 - \tilde{A}_0 \tilde{A}_3 > 0\}, \\ \mathcal{A}_4^+ &= \{\tilde{A}_0 > 0; \tilde{A}_1 > 0; \tilde{A}_3 > \tilde{A}_4 \tilde{A}_1^2 / (\tilde{A}_1 \tilde{A}_2 - \tilde{A}_0 \tilde{A}_3); \tilde{A}_1 \tilde{A}_2 - \tilde{A}_0 \tilde{A}_3 > 0, \tilde{A}_4 > 0\}.\end{aligned}$$

Notice that if we define

$$\mathcal{B} = \{\tilde{A}_0 > 0; \tilde{A}_1 > 0; \tilde{A}_3 > 0; \tilde{A}_1 \tilde{A}_2 - \tilde{A}_0 \tilde{A}_3 > 0; \tilde{A}_4 > 0\}$$

it is clear that $P(\mathcal{B}) = p_3^+/2$ and, moreover $\mathcal{A}_4^+ \subset \mathcal{B}$, with a strict inclusion. Hence $p_4^+ = P(\mathcal{A}_4^+) < P(\mathcal{B}) = p_3^+/2$, and $r_4 < r_3/2$, as we wanted to show. ■

Corollary 9. *Consider a linear random homogeneous differential equation of order $n = 3$ and the random variable X defined above. Then $p_0 = p_3 = 1/16$ and $p_1 = p_2 = 7/16$.*

Proof. By the above proposition, for $n = 3$, $p_0 = p_3 = r_3 = 1/16$. Hence, by Proposition 5, $p_1 = p_2 = 7/16$. ■

The computations in this case are similar to the ones of the previous section and the obtained results are summarized in Table 2. We only give some comments for the cases $n = 8$ and 10, where we have encountered that the vectors $\hat{\mathbf{p}}$ have negative and very small entries. This has occurred because the observed frequencies obtained by the Monte Carlo approach corresponding to these probabilities are not enough accurate. For this reason, we have made a new optimization step, again with the least squares method, but using as a vector to optimize the obtained $\hat{\mathbf{p}}$ vector but forcing those negative entries to be zero.

We explain this process for the $n = 8$ order case: The observed relative frequencies vector obtained by the Monte Carlo method is

$$\tilde{\mathbf{p}}^t = \left(\frac{1}{50000000}, \frac{6599}{50000000}, \frac{1159359}{50000000}, \frac{4996163}{20000000}, \frac{45377377}{100000000}, \frac{4995607}{20000000}, \frac{2318357}{100000000}, \frac{13497}{100000000}, \frac{1}{100000000} \right).$$

The relations stated in Corollary 6 are $p_3 = 1/4 - p_1$, $p_4 = 1/2 - 2p_0 - 2p_2$, $p_5 = p_3$, $p_6 = p_2$, $p_7 = p_1$, $p_8 = p_0$. By solving the system (7) with

$$M = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 0 \\ -2 & 0 & -2 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}; \quad \mathbf{q} = \begin{pmatrix} p_0 \\ p_1 \\ p_2 \end{pmatrix}; \quad \text{and } b = \begin{pmatrix} 0 \\ 0 \\ 0 \\ \frac{1}{2} \\ \frac{1}{4} \\ \frac{1}{2} \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

we obtain

$$\hat{\mathbf{q}}^t = \left(-\frac{5779}{200000000}, \frac{13569}{80000000}, \frac{4631293}{200000000} \right).$$

Hence, by (9) we get

$$\hat{\mathbf{p}} = \left(\frac{-5779}{200000000}, \frac{13569}{80000000}, \frac{4631293}{200000000}, \frac{19986431}{80000000}, \frac{22687243}{50000000}, \right. \\ \left. \frac{19986431}{80000000}, \frac{4631293}{200000000}, \frac{13569}{80000000}, \frac{-5779}{200000000} \right).$$

So we impose that $p_0 = p_8 = 0$. Thus we have $p_3 = p_5 = 1/4 - p_1$, $p_4 = 1/2 - 2p_0 - 2p_2 = 1/2 - 2p_2$, $p_6 = p_2$ and $p_7 = p_1$. We find the least squares solution of the system

$$\begin{pmatrix} \hat{p}_1 \\ \hat{p}_2 \\ \hat{p}_3 \\ \hat{p}_4 \\ \hat{p}_5 \\ \hat{p}_6 \\ \hat{p}_7 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \\ -1 & 0 \\ 0 & -2 \\ -1 & 0 \\ 0 & 1 \\ 1 & 0 \end{pmatrix} \cdot \begin{pmatrix} \hat{p}_1^* \\ \hat{p}_2^* \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ \frac{1}{4} \\ \frac{1}{2} \\ \frac{1}{4} \\ 0 \\ 0 \end{pmatrix}.$$

Using (8) and (9) we obtain

$$\hat{\mathbf{p}}^* = \left(0, \frac{13569}{80000000}, \frac{13882321}{600000000}, \frac{19986431}{80000000}, \frac{136117679}{300000000}, \frac{19986431}{80000000}, \frac{13882321}{600000000}, \frac{13569}{80000000}, 0 \right)$$

$$\simeq (0, 0.00017, 0.02314, 0.24983, 0.45373, 0.24983, 0.02314, 0.00017, 0).$$

The $n = 10$ case follows analogously.

Dimension	Observed frequency	Least squares	Relations (Corol. 6 and 9)
$n = 1$	$\tilde{p}_0 = 0.49997$ $\tilde{p}_1 = 0.50003$		$p_0 = 0.5$ $p_1 = 0.5$
$n = 2$	$\tilde{p}_0 = 0.24994$ $\tilde{p}_1 = 0.49999$ $\tilde{p}_2 = 0.25007$		$p_0 = 0.25$ $p_1 = 0.5$ $p_2 = 0.25$
$n = 3$	$\tilde{p}_0 = 0.06252$ $\tilde{p}_1 = 0.43743$ $\tilde{p}_2 = 0.43756$ $\tilde{p}_3 = 0.06249$		$p_0 = \frac{1}{16} = 0.0625$ $p_1 = \frac{7}{16} = 0.4375$ $p_2 = \frac{7}{16} = 0.4375$ $p_3 = \frac{1}{16} = 0.0625$
$n = 4$	$\tilde{p}_0 = 0.00928$ $\tilde{p}_1 = 0.24998$ $\tilde{p}_2 = 0.48152$ $\tilde{p}_3 = 0.24994$ $\tilde{p}_4 = 0.00929$	$\hat{p}_0 = 0.00925$ $\hat{p}_1 = 0.25$ $\hat{p}_2 = 0.48150$ $\hat{p}_3 = 0.25$ $\hat{p}_4 = 0.00925$	p_0 $p_1 = \frac{1}{4}$ $p_2 = \frac{1}{2} - 2p_0$ $p_3 = \frac{1}{4}$ $p_4 = p_0$
$n = 5$	$\tilde{p}_0 = 0.00071$ $\tilde{p}_1 = 0.08405$ $\tilde{p}_2 = 0.41526$ $\tilde{p}_3 = 0.41523$ $\tilde{p}_4 = 0.08404$ $\tilde{p}_5 = 0.00071$	$\hat{p}_0 = 0.00071$ $\hat{p}_1 = 0.08404$ $\hat{p}_2 = 0.41525$ $\hat{p}_3 = 0.41525$ $\hat{p}_4 = 0.08404$ $\hat{p}_5 = 0.00071$	p_0 p_1 $p_2 = \frac{1}{2} - p_0 - p_1$ $p_3 = \frac{1}{2} - p_0 - p_1$ $p_4 = p_1$ $p_5 = p_0$
$n = 6$	$\tilde{p}_0 = 0.00003$ $\tilde{p}_1 = 0.01723$ $\tilde{p}_2 = 0.24994$ $\tilde{p}_3 = 0.46562$ $\tilde{p}_4 = 0.24993$ $\tilde{p}_5 = 0.01723$ $\tilde{p}_6 = 0.00003$	$\hat{p}_0 = 0.00005$ $\hat{p}_1 = 0.01720$ $\hat{p}_2 = 0.24995$ $\hat{p}_3 = 0.46560$ $\hat{p}_4 = 0.24995$ $\hat{p}_5 = 0.01720$ $\hat{p}_6 = 0.00005$	p_0 p_1 $p_2 = \frac{1}{4} - p_0$ $p_3 = \frac{1}{2} - 2p_1$ $p_4 = \frac{1}{4} - p_0$ $p_5 = p_1$ $p_6 = p_0$
$n = 7$	$\tilde{p}_0 = 0.00000$ $\tilde{p}_1 = 0.00200$ $\tilde{p}_2 = 0.09571$ $\tilde{p}_3 = 0.40224$ $\tilde{p}_4 = 0.40233$ $\tilde{p}_5 = 0.09573$ $\tilde{p}_6 = 0.00199$ $\tilde{p}_7 = 0.00000$	$\hat{p}_0 = 0.00000$ $\hat{p}_1 = 0.00200$ $\hat{p}_2 = 0.09572$ $\hat{p}_3 = 0.40228$ $\hat{p}_4 = 0.40228$ $\hat{p}_5 = 0.09572$ $\hat{p}_6 = 0.00200$ $\hat{p}_7 = 0.00000$	p_0 p_1 p_2 $p_3 = \frac{1}{2} - p_0 - p_1 - p_2$ $p_4 = \frac{1}{2} - p_0 - p_1 - p_2$ $p_5 = p_2$ $p_6 = p_1$ $p_7 = p_0$

Dimension	Observed frequency	Least squares	Relations (Corol. 6 and 9)
$n = 8$	$\tilde{p}_0 = 0.00000$ $\tilde{p}_1 = 0.00013$ $\tilde{p}_2 = 0.02319$ $\tilde{p}_3 = 0.24981$ $\tilde{p}_4 = 0.45377$ $\tilde{p}_5 = 0.24978$ $\tilde{p}_6 = 0.02318$ $\tilde{p}_7 = 0.00013$ $\tilde{p}_8 = 0.00000$	$\hat{p}_0^* = 0.00000$ $\hat{p}_1^* = 0.00017$ $\hat{p}_2^* = 0.02314$ $\hat{p}_3^* = 0.24983$ $\hat{p}_4^* = 0.45372$ $\hat{p}_5^* = 0.24983$ $\hat{p}_6^* = 0.02314$ $\hat{p}_7^* = 0.00017$ $\hat{p}_8^* = 0.00000$	p_0 p_1 p_2 $p_3 = \frac{1}{4} - p_1$ $p_4 = \frac{1}{2} - 2p_0 - 2p_2$ $p_5 = \frac{1}{4} - p_1$ $p_6 = p_2$ $p_7 = p_1$ $p_8 = p_0$
$n = 9$	$\tilde{p}_0 = 0$ $\tilde{p}_1 = 0.00001$ $\tilde{p}_2 = 0.00336$ $\tilde{p}_3 = 0.10337$ $\tilde{p}_4 = 0.39328$ $\tilde{p}_5 = 0.39328$ $\tilde{p}_6 = 0.10332$ $\tilde{p}_7 = 0.00338$ $\tilde{p}_8 = 0.00000$ $\tilde{p}_9 = 0$	$\hat{p}_0 = 0$ $\hat{p}_1 = 0.00005$ $\hat{p}_2 = 0.00337$ $\hat{p}_3 = 0.10335$ $\hat{p}_4 = 0.39328$ $\hat{p}_5 = 0.39328$ $\hat{p}_6 = 0.10335$ $\hat{p}_7 = 0.00337$ $\hat{p}_8 = 0.00005$ $\hat{p}_9 = 0$	p_0 p_1 p_2 p_3 $p_4 = \frac{1}{2} - p_0 - p_1 - p_2 - p_3$ $p_5 = \frac{1}{2} - p_0 - p_1 - p_2 - p_3$ $p_6 = p_3$ $p_7 = p_2$ $p_8 = p_1$ $p_9 = p_0$
$n = 10$	$\tilde{p}_0 = 0$ $\tilde{p}_1 = 0.00000$ $\tilde{p}_2 = 0.00030$ $\tilde{p}_3 = 0.02784$ $\tilde{p}_4 = 0.24976$ $\tilde{p}_5 = 0.44421$ $\tilde{p}_6 = 0.24973$ $\tilde{p}_7 = 0.02787$ $\tilde{p}_8 = 0.00029$ $\tilde{p}_9 = 0.00000$ $\tilde{p}_{10} = 0$	$\hat{p}_0^* = 0.00000$ $\hat{p}_1^* = 0.00002$ $\hat{p}_2^* = 0.00028$ $\hat{p}_3^* = 0.02787$ $\hat{p}_4^* = 0.24972$ $\hat{p}_5^* = 0.44422$ $\hat{p}_6^* = 0.24972$ $\hat{p}_7^* = 0.02787$ $\hat{p}_8^* = 0.00028$ $\hat{p}_9^* = 0.00002$ $\hat{p}_{10}^* = 0.00000$	p_0 p_1 p_2 p_3 $p_4 = \frac{1}{4} - p_0 - p_2$ $p_5 = \frac{1}{2} - 2p_1 - 2p_3$ $p_6 = \frac{1}{4} - p_0 - p_2$ $p_7 = p_3$ $p_8 = p_2$ $p_9 = p_1$ $p_{10} = p_0$

Table 2. Stability indices for order n linear random homogeneous differential equations.

6 Linear random maps

In order to keep the approach of the preceding sections, we suggest to consider random linear discrete dynamical systems of the form

$$\mathcal{B} \mathbf{x}_{k+1} = A \mathbf{x}_k \text{ where } \mathbf{x} \in \mathbb{R}^n, \quad (13)$$

where \mathcal{B} and each of the n^2 entries of the random matrix A are i.i.d. $N(0, 1)$ random variables. Then, given a linear discrete random system (13), its characteristic random polynomial associated to the matrix $\frac{1}{\mathcal{B}}A$ is

$$Q(\lambda) = Q_n \lambda^n + Q_{n-1} \lambda^{n-1} + \dots + Q_1 \lambda + Q_0$$

where each random variable Q_j is a polynomial expression of $1/\mathcal{B}, A_{1,1}, \dots, A_{n,n}$ with not an easy distribution law. We denote by X the random variables that assigns to each Q its number of roots with modulus smaller than 1, that is, the stability index of the matrix $\frac{1}{\mathcal{B}}A$. Also p_k denotes the probabilities that X takes the value k .

As we will see in the examples, in this case the condition $p_k = p_{n-k}$ is no more satisfied. Among other reasons it happens that the entries of A^{-1} have not simple distributions. Since we do not know other relations on the probabilities p_k than the trivial one $\sum_{k=0}^n p_k = 1$, which is directly fulfilled by the observed relative frequencies, in this case we do not perform the least squares refinement.

The case $n = 1$ is the only one that we have been able to solve analytically. Notice that in this situation the only eigenvalue of $Q(\lambda)$ is $\lambda = A/B$, with A and B independent and $N(0, 1)$. Hence $p_0 = P(|A/B| > 1)$ and $p_1 = P(|A/B| < 1) = P(|B/A| > 1)$. Since A/B and B/A have the same distribution it holds that $p_0 = p_1 = 1/2$.

As in the other models, for each dimension $n \leq 10$, we generate 10^8 discrete systems of the form (13). For each matrix $\frac{1}{\mathcal{B}}A$ we have computed the characteristic polynomial Q and its associated polynomial Q^* (see Equation (6)) and have counted the number of roots of this last polynomial by using the Routh-Hurwitz zero counter. For $n \geq 5$ and in order to decrease the computation time we have directly numerically computed the eigenvalues of the matrix and counted the number of them with modulus less than one. The results obtained are shown in Table 3.

Dimension	Observed frequency
$n = 1$	$\tilde{p}_0 = 0.49994$ $\tilde{p}_1 = 0.50006$
$n = 2$	$\tilde{p}_0 = 0.46348$ $\tilde{p}_1 = 0.27705$ $\tilde{p}_2 = 0.25947$
$n = 3$	$\tilde{p}_0 = 0.45261$ $\tilde{p}_1 = 0.25828$ $\tilde{p}_2 = 0.15351$ $\tilde{p}_3 = 0.13560$
$n = 4$	$\tilde{p}_0 = 0.45040$ $\tilde{p}_1 = 0.24732$ $\tilde{p}_2 = 0.14799$ $\tilde{p}_3 = 0.08127$ $\tilde{p}_4 = 0.07302$
$n = 5$	$\tilde{p}_0 = 0.44957$ $\tilde{p}_1 = 0.24536$ $\tilde{p}_2 = 0.13956$ $\tilde{p}_3 = 0.08116$ $\tilde{p}_4 = 0.04443$ $\tilde{p}_5 = 0.03992$
$n = 6$	$\tilde{p}_0 = 0.44944$ $\tilde{p}_1 = 0.24419$ $\tilde{p}_2 = 0.13838$ $\tilde{p}_3 = 0.07536$ $\tilde{p}_4 = 0.04606$ $\tilde{p}_5 = 0.02449$ $\tilde{p}_6 = 0.02209$
$n = 7$	$\tilde{p}_0 = 0.44937$ $\tilde{p}_1 = 0.24394$ $\tilde{p}_2 = 0.13723$ $\tilde{p}_3 = 0.07480$ $\tilde{p}_4 = 0.04226$ $\tilde{p}_5 = 0.02636$ $\tilde{p}_6 = 0.01367$ $\tilde{p}_7 = 0.01236$

Dimension	Observed frequency
$n = 8$	$\tilde{p}_0 = 0.44937$ $\tilde{p}_1 = 0.24381$ $\tilde{p}_2 = 0.13702$ $\tilde{p}_3 = 0.07388$ $\tilde{p}_4 = 0.04207$ $\tilde{p}_5 = 0.02394$ $\tilde{p}_6 = 0.01526$ $\tilde{p}_7 = 0.00768$ $\tilde{p}_8 = 0.00698$
$n = 9$	$\tilde{p}_0 = 0.44941$ $\tilde{p}_1 = 0.24374$ $\tilde{p}_2 = 0.13680$ $\tilde{p}_3 = 0.07371$ $\tilde{p}_4 = 0.04139$ $\tilde{p}_5 = 0.02400$ $\tilde{p}_6 = 0.01374$ $\tilde{p}_7 = 0.00889$ $\tilde{p}_8 = 0.00434$ $\tilde{p}_9 = 0.00397$
$n = 10$	$\tilde{p}_0 = 0.44934$ $\tilde{p}_1 = 0.24371$ $\tilde{p}_2 = 0.13687$ $\tilde{p}_3 = 0.07358$ $\tilde{p}_4 = 0.04129$ $\tilde{p}_5 = 0.02348$ $\tilde{p}_6 = 0.01388$ $\tilde{p}_7 = 0.00792$ $\tilde{p}_8 = 0.00520$ $\tilde{p}_9 = 0.00247$ $\tilde{p}_{10} = 0.00226$

Table 3. Stability indices for linear random maps.

7 Linear random difference equations of order n

Finally we consider difference equations of order n of type

$$A_n x_{k+n} + A_{n-1} x_{k+n-1} + \cdots + A_1 x_{k+1} + A_0 x_k = 0,$$

where all the coefficients are i.i.d. random variables with $N(0, 1)$ distribution. In this situation, the stability index is given by the number of zeros with modulus smaller than 1 of the random characteristic polynomial $Q(\lambda) = A_n \lambda^n + A_{n-1} \lambda^{n-1} + \cdots + A_1 \lambda + A_0$. As in the preceding sections let X be the random variable that counts the number of roots of $Q(\lambda)$ with modulus smaller than 1 and set $p_k = P(X = k)$ for $k = 0, 1, \dots, n$.

Before proving some relations among the probabilities p_k , we give two preliminary lemmas. Let $\operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-u^2} du$ be the error function. The following result is stated in [3]. We prove it for the sake of completeness.

Lemma 10. *For $\alpha > 0$ and $\beta \in \mathbb{R}$,*

$$F(\alpha, \beta) := \int_0^\infty e^{-\alpha^2 x^2} \operatorname{erf}(\beta x) dx = \frac{\arctan(\beta/\alpha)}{\alpha\sqrt{\pi}}.$$

Proof. Fixed $\alpha > 0$, the function that defines F is absolutely integrable because $|\operatorname{erf}(x)| \leq 1$. Moreover its partial derivative with respect to β is also absolutely integrable. Hence $\lim_{\beta \rightarrow 0} F(\alpha, \beta) = F(\alpha, 0) = 0$ and

$$\begin{aligned} \frac{\partial F(\alpha, \beta)}{\partial \beta} &= \int_0^\infty \frac{\partial}{\partial \beta} \left(e^{-\alpha^2 x^2} \operatorname{erf}(\beta x) \right) dx = \frac{2}{\sqrt{\pi}} \int_0^\infty x e^{-\alpha^2 x^2} e^{-\beta^2 x^2} dx \\ &= \frac{2}{\sqrt{\pi}} \int_0^\infty x e^{-(\alpha^2 + \beta^2)x^2} dx = \frac{1}{(\alpha^2 + \beta^2)\sqrt{\pi}}. \end{aligned}$$

Therefore

$$F(\alpha, \beta) = F(\alpha, 0) + \int_0^\beta \frac{\partial F(\alpha, t)}{\partial t} dt = \int_0^\beta \frac{1}{(\alpha^2 + t^2)\sqrt{\pi}} dt = \frac{\arctan(\beta/\alpha)}{\alpha\sqrt{\pi}},$$

as we wanted to prove. ■

Next result is a consequence of the previous lemma.

Lemma 11. *Let $U \sim N(0, \sigma^2)$ and $V \sim N(0, \rho^2)$ be independent normal random variables. Then $P(U^2 - V^2 > 0) = \frac{2}{\pi} \arctan(\sigma/\rho)$.*

Proof. The joint density function of the random vector (U, V) is $f_\sigma(u)f_\rho(v)$, where $f_s(u) = e^{-u^2/(2s^2)}/(\sqrt{2\pi}s)$. Observe that the points $(u, v) \in \mathbb{R}^2$ such that $u^2 - v^2 > 0$ is the region

where $-|u| < v < |u|$, hence by symmetry,

$$\begin{aligned} P(U^2 - V^2 > 0) &= 4 \int_0^\infty f_\sigma(u) \int_0^u f_\rho(v) dv du = \frac{4}{2\pi\sigma\rho} \int_0^\infty e^{-u^2/(2\sigma^2)} \int_0^u e^{-v^2/(2\rho^2)} dv du \\ &= \frac{2}{\pi\sigma\rho} \int_0^\infty e^{-u^2/(2\sigma^2)} \operatorname{erf}\left(\frac{u}{\sqrt{2}\rho}\right) \sqrt{\frac{\pi}{2}}\rho du = \frac{2}{\pi} \arctan\left(\frac{\sigma}{\rho}\right), \end{aligned}$$

where in the last equality we have used Lemma 10. ■

Notice that with the notations of the above lemma, $P(U^2 - V^2 > 0) + P(U^2 - V^2 < 0) = 1$. Hence

$$P(U^2 - V^2 < 0) = 1 - \frac{2}{\pi} \arctan\left(\frac{\sigma}{\rho}\right) = \frac{2}{\pi} \arctan\left(\frac{\rho}{\sigma}\right), \quad (14)$$

where we have used that $\arctan(x) + \arctan(1/x) = \pi/2$ or, simply, the same lemma interchanging U and V . Observe also that when $\sigma = \rho$, $P(U^2 - V^2 > 0) = P(U^2 - V^2 < 0) = 1/2$, result that, in fact, is a straightforward consequence that in this situation $U^2 - V^2$ and $V^2 - U^2$ have the same distribution.

Proposition 12. *With the above notations:*

- (a) $\sum_{k=0}^n p_k = 1$.
- (b) For all $k \in \{0, 1, \dots, n\}$, $p_k = p_{n-k}$.
- (c) When n is odd, $\sum_{i \text{ even}} p_i = \sum_{i \text{ odd}} p_i = \frac{1}{2}$.
- (d) When $n = 2k$ is even,

$$\sum_{i \text{ even}} p_i = \frac{2}{\pi} \arctan\left(\sqrt{\frac{k+1}{k}}\right) \quad \text{and} \quad \sum_{i \text{ odd}} p_i = \frac{2}{\pi} \arctan\left(\sqrt{\frac{k}{k+1}}\right). \quad (15)$$

Proof. The first assert is obvious. To see the second one we compare the difference equation $a_n x_{k+n} + a_{n-1} x_{k+n-1} + \dots + a_1 x_{k+1} + a_0 x_k = 0$, with $a_i \in \mathbb{R}$, $i = 0, 1, \dots, n$, with characteristic polynomial $Q(\lambda) = a_n \lambda^n + a_{n-1} \lambda^{n-1} + \dots + a_2 \lambda^2 + a_1 \lambda + a_0$ with the difference equation $a_n x_k + a_{n-1} x_{k+1} + \dots + a_0 x_{k+n} = 0$ with characteristic polynomial $Q^*(\lambda) = a_n + a_{n-1} \lambda + \dots + a_1 \lambda^{n-1} + a_0 \lambda^n$. Notice that if $Q(\lambda)$ has m non-zero roots with modulus smaller than 1 and $n - m$ with modulus bigger than 1, then the converse follows for $Q^*(\lambda)$ because $Q(\lambda) = 0$ if and only if $Q^*(\frac{1}{\lambda}) = 0$. From this result applied to the corresponding random polynomials we get that $p_k = p_{n-k}$, because both have identically distributed coefficients. So we have proved statement (b). To prove items (c) and (d) recall first that it was proved in item (c) of Proposition 5 that a polynomial, with no roots with

zero real part, $Q(\lambda) = a_n\lambda^n + a_{n-1}\lambda^{n-1} + \dots + a_2\lambda^2 + a_1\lambda + a_0$ has an even number of roots with negative real part if and only if $a_n a_0 > 0$. By using the polynomial

$$\begin{aligned} Q^*(z) &= a_n(z+1)^n + a_{n-1}(z+1)^{n-1}(z-1) + \dots + a_0(z-1)^n \\ &= (a_n + a_{n-1} + \dots + a_1 + a_0)z^n + \dots + (a_n - a_{n-1} + a_{n-2} - \dots + (-1)^n a_0), \end{aligned}$$

introduced in Section 3 (Equation (6)) we get that $Q(\lambda)$, with no roots of modulus 1, has an even number ($2m$) of roots with modulus smaller than 1 if and only if $Q^*(z)$ has exactly $2m$ roots with negative real part and this happens if and only if $(a_n + a_{n-1} + \dots + a_1 + a_0) \cdot (a_n - a_{n-1} + a_{n-2} - \dots + (-1)^n a_0) > 0$. Hence, considering the corresponding random polynomials, we have that

$$\begin{aligned} \sum_{i \text{ even}} p_i &= P((A_n + A_{n-1} + \dots + A_0) \cdot (A_n - A_{n-1} + \dots + (-1)^n A_0) > 0) \\ &= P(U^2 - V^2 > 0), \end{aligned}$$

where $U = A_n + A_{n-2} + A_{n-4} + \dots$ and $V = A_{n-1} + A_{n-3} + A_{n-5} + \dots$ and the sums end either at A_0 or A_1 according the parity of n . Since $A_j \sim N(0, 1)$ and all A_j are independent we get that when $n = 2k$ (resp. $n = 2k-1$) we have that $U \sim N(0, k+1)$ (resp. $U \sim N(0, k)$) and $V \sim N(0, k)$ and U and V are independent. Hence, by using Lemma 11, we obtain that when $n = 2k - 1$, $P(U^2 - V^2 > 0) = 1/2$ and that when $n = 2k$,

$$\sum_{i \text{ even}} p_i = P(U^2 - V^2 > 0) = \frac{2}{\pi} \arctan\left(\sqrt{\frac{k+1}{k}}\right).$$

The sum of all p_i when i is odd can be obtained from the previous one, see also (14). ■

Corollary 13. (i) Consider the linear random homogeneous difference equation of order n , let X be the random variable defined above and $p_k = P(X = k)$. Then the probabilities p_k satisfy all the consequences of Lemma 2. In particular $E(X) = n/2$.

(ii) Moreover the new affine relations given in Equations (15) hold. In particular, for $n = 2$, $p_0 = p_2 = \frac{1}{\pi} \arctan(\sqrt{2})$ and $p_1 = \frac{2}{\pi} \arctan(\frac{1}{\sqrt{2}})$; and for $n = 4$, $p_1 = p_3 = \frac{1}{\pi} \arctan(\sqrt{\frac{2}{3}})$.

In this case, and for the situations where we have not been able to obtain the exact probabilities we have done similar computations that in the previous section, first with the Monte Carlo method, generating for each order $n = 0, \dots, 10$, 10^8 random vectors $(A_0, \dots, A_n) \in \mathbb{R}^{n+1}$ whose components are pseudo-random numbers with $N(0, 1)$ distribution. Then, by using the relations in Proposition 12 and Corollary 13 we have performed a least squares refinement.

For instance for $n = 4$, by Corollary 13 we have $p_1 = p_3 = \arctan(\sqrt{2/3})/\pi \simeq 0.217953$; $p_2 = 2 \arctan(\sqrt{3/2})/\pi - 2p_0$ and $p_4 = p_0$. Hence, we fix the values $\hat{p}_1 = p_1$ and $\hat{p}_3 = p_3$ and system (7) writes

$$M\hat{\mathbf{q}} + b = \begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix} \cdot (\hat{p}_0) + \begin{pmatrix} 0 \\ \frac{2}{\pi} \arctan\left(\sqrt{\frac{3}{2}}\right) \\ 0 \end{pmatrix} = \begin{pmatrix} \tilde{p}_0 \\ \tilde{p}_2 \\ \tilde{p}_4 \end{pmatrix}.$$

Hence we can easily find the least squares solution of the above incompatible linear system:

$$(1, -2, 1) \cdot \left[\begin{pmatrix} 1 \\ -2 \\ 1 \end{pmatrix} \cdot (\hat{p}_0) + \begin{pmatrix} 0 \\ \frac{2}{\pi} \arctan\left(\sqrt{\frac{3}{2}}\right) \\ 0 \end{pmatrix} \right] = (1, -2, 1) \cdot \begin{pmatrix} \tilde{p}_0 \\ \tilde{p}_2 \\ \tilde{p}_4 \end{pmatrix},$$

thus we get the result in Equation (8): $\hat{p}_0 = \frac{1}{6}(\tilde{p}_0 - 2\tilde{p}_2 + \tilde{p}_4) + \frac{2}{3\pi} \arctan\left(\sqrt{3/2}\right)$, and therefore $\hat{p}_4 = \hat{p}_0$ and $\hat{p}_2 = 2 \arctan(\sqrt{3/2})/\pi - 2\hat{p}_0$. Since our Monte Carlo simulations give

$$(\tilde{p}_0, \tilde{p}_2, \tilde{p}_4) = \left(\frac{2056203}{20000000}, \frac{7169499}{20000000}, \frac{10285619}{100000000} \right) \simeq (0.10281, 0.35847, 0.10286),$$

the above relations show that $(\hat{p}_0, \hat{p}_2, \hat{p}_4) \simeq (0.10282, 0.35846, 0.10282)$.

All our results are collected in Table 4.

Order	Observed frequency	Least squares	Relations (Prop. 12 and Cor. 13))
$n = 1$	$\tilde{p}_0 = 0.49991$ $\tilde{p}_1 = 0.50009$		$p_0 = 0.5$ $p_1 = 0.5$
$n = 2$	$\tilde{p}_0 = 0.30410$ $\tilde{p}_1 = 0.39184$ $\tilde{p}_2 = 0.30406$		$p_0 = \frac{1}{\pi} \arctan(\sqrt{2}) \simeq 0.304087$ $p_1 = \frac{2}{\pi} \arctan\left(\frac{1}{\sqrt{2}}\right) \simeq 0.391826$ $p_2 = \frac{1}{\pi} \arctan(\sqrt{2}) \simeq 0.304087$
$n = 3$	$\tilde{p}_0 = 0.17251$ $\tilde{p}_1 = 0.32752$ $\tilde{p}_2 = 0.32753$ $\tilde{p}_3 = 0.17244$	$\hat{p}_0 = 0.17248$ $\hat{p}_1 = 0.32752$ $\hat{p}_2 = 0.32752$ $\hat{p}_3 = 0.17248$	p_0 $p_1 = \frac{1}{2} - p_0$ $p_2 = \frac{1}{2} - p_0$ $p_3 = p_0$
$n = 4$	$\tilde{p}_0 = 0.10281$ $\tilde{p}_1 = 0.21792$ $\tilde{p}_2 = 0.35847$ $\tilde{p}_3 = 0.21794$ $\tilde{p}_4 = 0.10286$	$\hat{p}_0 = 0.10282$ $\hat{p}_1 = 0.21795$ $\hat{p}_2 = 0.35846$ $\hat{p}_3 = 0.21795$ $\hat{p}_4 = 0.10282$	p_0 $p_1 = \frac{1}{\pi} \arctan\left(\sqrt{\frac{2}{3}}\right) \simeq 0.217953$ $p_2 = \frac{2}{\pi} \arctan\left(\sqrt{\frac{3}{2}}\right) - 2p_0$ $p_3 = \frac{1}{\pi} \arctan\left(\sqrt{\frac{2}{3}}\right) \simeq 0.217953$ $p_4 = p_0$

Order	Observed frequency	Least squares	Relations (Prop. 12 and Cor. 13)
$n = 5$	$\tilde{p}_0 = 0.05909$ $\tilde{p}_1 = 0.15331$ $\tilde{p}_2 = 0.28760$ $\tilde{p}_3 = 0.28756$ $\tilde{p}_4 = 0.15335$ $\tilde{p}_5 = 0.05908$	$\hat{p}_0 = 0.05909$ $\hat{p}_1 = 0.15333$ $\hat{p}_2 = 0.28758$ $\hat{p}_3 = 0.28758$ $\hat{p}_4 = 0.15333$ $\hat{p}_5 = 0.05909$	p_0 p_1 $p_2 = \frac{1}{2} - p_0 - p_1$ $p_3 = \frac{1}{2} - p_0 - p_1$ $p_4 = p_1$ $p_5 = p_0$
$n = 6$	$\tilde{p}_0 = 0.03501$ $\tilde{p}_1 = 0.09726$ $\tilde{p}_2 = 0.23777$ $\tilde{p}_3 = 0.25985$ $\tilde{p}_4 = 0.23781$ $\tilde{p}_5 = 0.09724$ $\tilde{p}_6 = 0.03505$	$\hat{p}_0 = 0.03502$ $\hat{p}_1 = 0.09726$ $\hat{p}_2 = 0.23779$ $\hat{p}_3 = 0.25986$ $\hat{p}_4 = 0.23779$ $\hat{p}_5 = 0.09726$ $\hat{p}_6 = 0.03502$	p_0 p_1 $p_2 = \frac{1}{\pi} \arctan(\sqrt{\frac{4}{3}}) - p_0$ $p_3 = \frac{2}{\pi} \arctan(\sqrt{\frac{3}{4}}) - 2p_1$ $p_4 = \frac{1}{\pi} \arctan(\sqrt{\frac{4}{3}}) - p_0$ $p_5 = p_1$ $p_6 = p_0$
$n = 7$	$\tilde{p}_0 = 0.02025$ $\tilde{p}_1 = 0.06432$ $\tilde{p}_2 = 0.17174$ $\tilde{p}_3 = 0.24376$ $\tilde{p}_4 = 0.24361$ $\tilde{p}_5 = 0.17177$ $\tilde{p}_6 = 0.06428$ $\tilde{p}_7 = 0.02025$	$\hat{p}_0 = 0.02025$ $\hat{p}_1 = 0.06430$ $\hat{p}_2 = 0.17176$ $\hat{p}_3 = 0.24369$ $\hat{p}_4 = 0.24369$ $\hat{p}_5 = 0.17176$ $\hat{p}_6 = 0.06430$ $\hat{p}_7 = 0.02025$	p_0 p_1 p_2 $p_3 = \frac{1}{2} - p_0 - p_1 - p_2$ $p_4 = \frac{1}{2} - p_0 - p_1 - p_2$ $p_5 = p_2$ $p_6 = p_1$ $p_7 = p_0$
$n = 8$	$\tilde{p}_0 = 0.01194$ $\tilde{p}_1 = 0.03994$ $\tilde{p}_2 = 0.12272$ $\tilde{p}_3 = 0.19230$ $\tilde{p}_4 = 0.25701$ $\tilde{p}_5 = 0.19238$ $\tilde{p}_6 = 0.12724$ $\tilde{p}_7 = 0.03994$ $\tilde{p}_8 = 0.01197$	$\hat{p}_0 = 0.01196$ $\hat{p}_1 = 0.03994$ $\hat{p}_2 = 0.12726$ $\hat{p}_3 = 0.19234$ $\hat{p}_4 = 0.25700$ $\hat{p}_5 = 0.19234$ $\hat{p}_6 = 0.12726$ $\hat{p}_7 = 0.03994$ $\hat{p}_8 = 0.01196$	p_0 p_1 p_2 $p_3 = \frac{1}{\pi} \arctan(\sqrt{\frac{4}{5}}) - p_1$ $p_4 = \frac{2}{\pi} \arctan(\sqrt{\frac{5}{4}}) - 2p_0 - 2p_2$ $p_5 = \frac{1}{\pi} \arctan(\sqrt{\frac{4}{5}}) - p_1$ $p_6 = p_2$ $p_7 = p_1$ $p_8 = p_0$

Order	Observed frequency	Least squares	Relations (Prop. 12 and Cor. 13)
$n = 9$	$\tilde{p}_0 = 0.00693$	$\hat{p}_0 = 0.00693$	p_0
	$\tilde{p}_1 = 0.02556$	$\hat{p}_1 = 0.02556$	p_1
	$\tilde{p}_2 = 0.08711$	$\hat{p}_2 = 0.08711$	p_2
	$\tilde{p}_3 = 0.15653$	$\hat{p}_3 = 0.15653$	p_3
	$\tilde{p}_4 = 0.22389$	$\hat{p}_4 = 0.22386$	$p_4 = \frac{1}{2} - p_0 - p_1 - p_2 - p_3$
	$\tilde{p}_5 = 0.22382$	$\hat{p}_5 = 0.22386$	$p_5 = \frac{1}{2} - p_0 - p_1 - p_2 - p_3$
	$\tilde{p}_6 = 0.15654$	$\hat{p}_6 = 0.15653$	$p_6 = p_3$
	$\tilde{p}_7 = 0.08712$	$\hat{p}_7 = 0.08711$	$p_7 = p_2$
	$\tilde{p}_8 = 0.02557$	$\hat{p}_8 = 0.02556$	$p_8 = p_1$
	$\tilde{p}_9 = 0.00693$	$\hat{p}_9 = 0.00693$	$p_9 = p_0$
$n = 10$	$\tilde{p}_0 = 0.00409$	$\hat{p}_0 = 0.00411$	p_0
	$\tilde{p}_1 = 0.01567$	$\hat{p}_1 = 0.01566$	p_1
	$\tilde{p}_2 = 0.06089$	$\hat{p}_2 = 0.06091$	p_2
	$\tilde{p}_3 = 0.11500$	$\hat{p}_3 = 0.11497$	p_3
	$\tilde{p}_4 = 0.19950$	$\hat{p}_4 = 0.19947$	$p_4 = \frac{1}{\pi} \arctan(\sqrt{\frac{6}{5}}) - p_0 - p_2$
	$\tilde{p}_5 = 0.20978$	$\hat{p}_5 = 0.20976$	$p_5 = \frac{2}{\pi} \arctan(\sqrt{\frac{5}{6}}) - 2p_1 - 2p_3$
	$\tilde{p}_6 = 0.19941$	$\hat{p}_6 = 0.19947$	$p_6 = \frac{1}{\pi} \arctan(\sqrt{\frac{6}{5}}) - p_0 - p_2$
	$\tilde{p}_7 = 0.11499$	$\hat{p}_7 = 0.11497$	$p_7 = p_3$
	$\tilde{p}_8 = 0.06088$	$\hat{p}_8 = 0.06091$	$p_8 = p_2$
	$\tilde{p}_9 = 0.01570$	$\hat{p}_9 = 0.01566$	$p_9 = p_1$
		$\tilde{p}_{10} = 0.00408$	$\hat{p}_{10} = 0.00411$

Table 4. Stability indices for order n linear random homogeneous difference equations.

References

- [1] J.C. Artés, J. Llibre. *Statistical measure of quadratic vector fields*. Resenhas 6 (2003), 85–97.
- [2] P. Bratley, B.L. Fox, L. E. Schrage. *A guide to simulation*, 2d ed. Springer, New York, 1987.
- [3] K. Briggs. *Integrals involving erf*. <http://keithbriggs.info/documents/erf-integrals.pdf>. (Accessed March 20, 2019).
- [4] B.M. Chen-Charpentier, D. Stanescu. *Epidemic models with random coefficients*. Math. Comput. Modelling 52 (2010), 1004–1010.

- [5] B.M. Chen-Charpentier, D. Stanescu. *Virus propagation with randomness*. Math. Comput. Modelling 57 (2013), 1816–1821
- [6] K. Conrad. *Probability distributions and maximum entropy*.
<http://www.math.uconn.edu/~kconrad/blurbs/analysis/entropypost.pdf> (Accessed November 11, 2018).
- [7] G. Dahlquist, Å. Björck. Numerical methods. Prentice-Hall, Englewood Cliffs NJ, 1974. Section 5.7 pp. 196–201.
- [8] I.S. Gradshteyn, I.M. Ryzhik, *Routh-Hurwitz Theorem*. §15.715 in Tables of Integrals, Series, and Products, 6th ed. Academic Press, San Diego CA, 2000, p. 1076.
- [9] M. Grigoriu, T. Soong. Random vibration in mechanical and structural systems. Prentice Hall, New Jersey, 1993.
- [10] F.B. Jones. *Connected and disconnected plane sets and the functional equation $f(x) + f(y) = f(x + y)$* . Bull. Amer. Math. Soc. 48 (1942), 115–120.
- [11] H. Lopes, B. Pagnoncelli, C. Palmeira. *Coeficientes aleatórios de equações diferenciais ordinárias lineares*. Matemática Universitária 14 (2008), 44–50. Translated to English at <http://bernardokp.uai.cl/> with title *Random linear systems and simulation*.
- [12] G. Marsaglia. *Choosing a point from the surface of a sphere*. The Annals of Mathematical Statistics 43 (1972), 645–646.
- [13] B.J.T. Morgan. Applied stochastic modelling. Arnold Publishers, London, 2000.
- [14] M.E. Muller. *A note on a method for generationg points uniformly on N-dimensional spheres*. Communications of the ACM 2 (1959), 19–20.
- [15] J.R. Schott. Matrix analysis for statistics. John Wiley & Sons, New York, 1997.
- [16] D. Stanescu, B.M. Chen-Charpentier. *Random coefficient differential equation models for bacterial growth*. Math. Comput. Modelling 50 (2009), 885–895.
- [17] J. Stoer, R. Bulirsch. Introduction to numerical analysis. Springer, New york, 1983.
- [18] S.H. Strogatz. Nonlinear dynamics and chaos. Westview Press, Cambridge MA, 1994.