

Around some extensions of Casas-Alvero conjecture for non-polynomial functions

A. Cima, A. Gasull, and F. Mañosas

Departament de Matemàtiques

Universitat Autònoma de Barcelona, Barcelona, Spain

`{cima,gasull,manyosas}@mat.uab.cat`

Abstract. We show that two natural extensions of the real Casas-Alvero conjecture in the non-polynomial setting do not hold.

2010 MSC: Primary: 30C15. Secondary: 12D10; 13P15; 26C10.

Keywords: *polynomial, Casas-Alvero conjecture, zeroes of functions*

1 Introduction and main results

The Casas-Alvero conjecture affirms that if a complex polynomial P of degree $n > 1$ shares roots with all its derivatives, $P^{(k)}$, $k = 1, 2, \dots, n-1$, then there exist two complex numbers, a and $b \neq 0$, such that $P(z) = b(z - a)^n$. Notice that, in principle, the common root between P and each $P^{(k)}$ might depend on k . Casas-Alvero arrived to this problem at the turn of this century, when he was working in his paper [1] trying to obtain an irreducibility criterion for two variable power series with complex coefficients. See [2] for an explanation of the problem in his own words.

Although several authors have got partial answers, to the best of our knowledge the conjecture remains open. For $n \leq 4$ the conjecture is a simple consequence of the wonderful Gauss-Lucas Theorem ([6]). In 2006 it was proved in [5], by using Maple, that it is true for $n \leq 8$. Afterwards in [6, 7] it was proved that it holds when n is $p^m, 2p^m, 3p^m$ or $4p^m$, for some prime number p and $m \in \mathbb{N}$. The first cases left open are those where $n = 24, 28$ or 30 . See again [6] for a very interesting survey or [3, 8] for some recent contributions on this question.

Adding the hypotheses that P is a real polynomial and all its n roots, taking into account their multiplicities, are real, the conjecture has a real counterpart, that also remains open. It says that $P(x) = b(x - a)^n$ for some real numbers a and $b \neq 0$. For this real case, the conjecture can be proved easily for $n \leq 4$, simply by using Rolle's Theorem. This tool does not suffice for $n \geq 5$, see for instance [4] for more details, or next section.

Also in the real case, in [6] it is proved that if the condition for one of the derivatives of P is removed, then there exist polynomials satisfying the remaining $n - 2$ conditions, different from $b(x - a)^n$. The construction of some of these polynomials presented in that paper is very nice and is a consequence of the Brouwer's fixed point Theorem in a suitable context.

Finally, it is known that if the conjecture holds in \mathbb{C} , then it is true over all fields of characteristic 0. On the other hand, it is not true over all fields of characteristic p , see again [7]. For instance, consider $P(x) = x^2(x^2 + 1)$ in characteristic 5 with roots 0, 0, 2 and 3. Then $P'(x) = 2x(2x^2 + 1)$, $P''(x) = 12x^2 + 2 = 2(x^2 + 1)$ and $P'''(x) = 4x$ and all them share roots with P .

The aim of this note is to present two natural extensions of the real Casas-Alvero conjecture to smooth functions and show that none of them holds.

Question 1. Fix $1 < n \in \mathbb{N}$. Let F be a class \mathcal{C}^n real function such that $F^{(n)}(x) \neq 0$ for all $x \in \mathbb{R}$, and has n real zeroes, taking into account their multiplicities. Assume that F shares zeroes with all its derivatives, $F^{(k)}$, $k = 1, 2, \dots, n - 1$. Is it true that $F(x) = b(f(x))^n$ for some $0 \neq b \in \mathbb{R}$ and some f , a class \mathcal{C}^n real function, that has exactly one simple zero?

Notice that one of the hypotheses of the real Casas-Alvero conjecture can be reformulated as follows: The polynomial F shares roots with all its derivatives but one, precisely the one corresponding to its degree. Trivially, this is so, because all the derivatives of order higher than n are identically zero. The second question that we consider is:

Question 2. Fix $1 < n \in \mathbb{N}$. Let F be a real analytic function that shares zeroes with all its derivatives but one, say $F^{(n)}$. Is it true that $F(x) = b(f(x))^n$ for some $0 \neq b \in \mathbb{R}$ and some real analytic function f , that has exactly one simple zero?

Theorem A. (i) The answer to the Question 1 is “yes” for $n \leq 4$ and “no” for $n = 5$.

(ii) The answer to the Question 2 is already “no” for $n = 2$.

Our result reinforces the intuitive idea that Casas-Alvero conjecture is mainly a question related with the rigid structure of the polynomials.

2 Proof of Theorem A

(i) The answer to Question 1 is “yes” for $n = 2, 3, 4$ because the proof of the real Casas-Alvero conjecture for the same values of n , based on the Rolle’s Theorem and given in [4], does not uses at all that P is a polynomial. Let us adapt it to our setting. Since $F^{(n)}$ does not vanish we know that F has exactly n real zeroes, taking into account their multiplicities. Moreover we know that F has to have at least a double zero, that without loss of generality can be taken as 0. Next we can do a case by case study to discard all situations except that F has only a zero and it is of multiplicity n . For the sake of brevity, we give the details only in one situation. Assume, to arrive to a contradiction, that $n = 4$, F is under the hypotheses of Question 1 and that its four zeroes are 0, 0, c_1, c_2 with $0 < c_1 < c_2$. By Rolle’s Theorem the zeroes of F' are 0, d_1 and d_2 and satisfy $0 < d_1 < c_1 < d_2 < c_2$. Then, since F'' has to have two zeroes, say e_1, e_2 , and they satisfy $0 < e_1 < d_1 < e_2 < d_2$, the hypotheses force that $e_2 = c_1$. Hence the zero of F''' has to be between e_1 and $e_2 = c_1$, that is in particular in $(0, c_1)$, interval that contains no zero of F , arriving to the desired contradiction. In short, we have proved for $n \leq 4$, that $F(x) = x^n G(x)$, for some class \mathcal{C}^n function G , that does not vanish. Hence

$$F(x) = \text{sign}(G(0)) \left(x \sqrt[n]{\frac{G(x)}{\text{sign}(G(0))}} \right)^n = b(f(x))^n,$$

where f has only one zero, $x = 0$, that is simple, as we wanted to prove.

To find a map F for which the answer to Question 1 is “no” we consider $n = 5$ and a configuration of zeroes of F and its derivatives proposed in [4] as the simplest one, compatible with the hypotheses of the Casas-Alvero conjecture and Rolle’s Theorem. Specifically, we will search for a function F , of class at least C^5 , with the five zeroes $0, 0, 1, c, d$, to be fixed, satisfying $0 < 1 < c < d$, and moreover

$$F'(0) = 0, \quad F''(1) = 0, \quad F'''(c) = 0, \quad F^{(4)}(1) = 0, \quad (1)$$

and such that $F^{(5)}$ does not vanish. Notice that $F'(0) = 0$ is not a new restriction.

We start assuming that $F^{(5)}(x) = r - \sin(x)$, for some $r > 1$ to be determined. By imposing that conditions (1) hold, together with $F(0) = 0$, we get that

$$F(x) = \int_0^x \int_0^u \int_1^w \int_c^z \int_1^y (r - \sin(t)) dt dy dz dw du.$$

Some straightforward computations give that

$$\begin{aligned} F(x) = & \frac{r}{120}x^5 - \frac{r + \cos(1)}{12}x^4 + \frac{2rc - 2\sin(c) + 2\cos(1)c - rc^2}{12}x^3 \\ & + \frac{6\sin(c) + 2r + 9\cos(1) - 6rc + 3rc^2 - 6\cos(1)c}{12}x^2 - 1 + \cos(x). \end{aligned}$$

Imposing now that $F(1) = 0$ we obtain that

$$r = \frac{5(8\cos(1)c - 41\cos(1) - 8\sin(c) + 24)}{4(5c^2 - 10c + 4)} = R(c).$$

Next we have to impose that $F(c) = 0$. By replacing the above expression of r in F we obtain that

$$F(c) = \frac{G(c)}{96(5c^2 - 10c + 4)},$$

where

$$\begin{aligned} G(c) = & -c^2(12c^4 - 369c^3 + 1437c^2 - 1708c + 532)\cos(1) - 8c^2(c-1)(c-2)^2\sin(c) \\ & + (480c^2 - 960c + 384)\cos(c) - 24(c-1)(9c^4 - 36c^3 + 24c^2 + 24c - 16). \end{aligned}$$

A carefully study shows that G has exactly one real zero $c_1 \in (17/10, 19/10) = I$, with $c_1 \approx 1.79343096$. To prove its existence it suffices to show that

$$\begin{aligned} G\left(\frac{17}{10}\right) &= -\frac{99211099}{500000}\cos(1) - \frac{18207}{12500}\sin\left(\frac{17}{10}\right) + \frac{696}{5}\cos\left(\frac{17}{10}\right) + \frac{1583211}{12500} > 0, \\ G\left(\frac{19}{10}\right) &= -\frac{180110481}{500000}\cos(1) - \frac{3249}{12500}\sin\left(\frac{19}{10}\right) + \frac{1464}{5}\cos\left(\frac{19}{10}\right) + \frac{3616677}{12500} < 0. \end{aligned}$$

By using Taylor’s formula we know that for any $c > 0$, $S^-(c) < \sin(c) < S^+(c)$ and $C^-(c) < \cos(c) < C^+(c)$ where

$$S^\pm(c) = c - \frac{c^3}{3!} + \frac{c^5}{5!} - \frac{c^7}{7!} + \frac{c^9}{9!} \pm \frac{c^{11}}{11!} \quad \text{and} \quad C^\pm(c) = 1 - \frac{c^2}{2!} + \frac{c^4}{4!} - \frac{c^6}{6!} + \frac{c^8}{8!} \pm \frac{c^{10}}{10!}.$$




Figure 1: Plot of a map F for which the answer to Question 1 for $n = 5$ is “no”.

Hence we can replace the values of the trigonometric functions in G by rational numbers to have upper or lower bounds of this function evaluated at 1, $17/10$ or $19/10$. For instance,

$$0.5403023 \approx \frac{1960649}{3628800} = C^-(1) < \cos(1) < C^+(1) = \frac{280093}{518400} \approx 0.5403028.$$

We obtain that

$$\begin{aligned} G\left(\frac{17}{10}\right) &> -\frac{99211099}{500000} C^+(1) - \frac{18207}{12500} S^+\left(\frac{17}{10}\right) + \frac{696}{5} C^-\left(\frac{17}{10}\right) + \frac{1583211}{12500} \\ &= \frac{3444600099561969856969}{4989600000000000000000} > 0 \end{aligned}$$

and

$$\begin{aligned} G\left(\frac{19}{10}\right) &< -\frac{180110481}{500000} C^-(1) - \frac{3249}{12500} S^-\left(\frac{19}{10}\right) + \frac{1464}{5} C^+\left(\frac{19}{10}\right) + \frac{3616677}{12500} \\ &= -\frac{1689627895469649855823}{1663200000000000000000} < 0. \end{aligned}$$

To show the uniqueness of the zero in I , we will prove that G is decreasing in this interval. It holds that

$$G'(c) = T(c) \cos(1) + U(c) \sin(c) + V(c) \cos(c) + W(c),$$

with

$$\begin{aligned} U(c) &= -8(5c^2 - 10c + 4)(c^2 - 2c + 12), & V(c) &= -8(c-1)(c^4 - 4c^3 + 4c^2 - 120), \\ T(c) &= -c(72c^4 - 1845c^3 + 5748c^2 - 5124c + 1064), & W(c) &= -120(9c^4 - 36c^3 + 36c^2 - 8). \end{aligned}$$

By computing the Sturm sequences of T, U and V we can prove that $T(c) < 0$, $U(c) < 0$ and $V(c) > 0$ for all $c \in I$. Hence, for $c \in I$,

$$G'(c) < T(c)C^-(c) + U(c)S^-(c) + V(c)C^+(c) + W(c) = Q(c),$$

where

$$Q(c) = \frac{72469}{64800}c - \frac{669211}{43200}c^2 + \frac{18852329}{302400}c^3 - \frac{8854991}{80640}c^4 + \frac{4732471}{50400}c^5 - \frac{532}{15}c^6 + \frac{8}{7}c^7 + \frac{191}{70}c^8 \\ - \frac{797}{1890}c^9 - \frac{34}{405}c^{10} + \frac{1651}{103950}c^{11} + \frac{3533}{2494800}c^{12} - \frac{193}{623700}c^{13} + \frac{1}{142560}c^{14} - \frac{1}{831600}c^{15}.$$

The Sturm sequence of Q shows that it has no zeroes in I . Moreover, it is negative in this interval, and as a consequence, G' is also negative, as we wanted to prove.

We fix $c = c_1$. Then, $r = R(c_1)$ and F is also totally fixed. Moreover, by using the same techniques we get that $r = R(c_1) > R(19/10) > 1$ and as a consequence $F^{(5)}$ does not vanish. In fact, $r = R(c_1) \approx 1.04591089$. Finally, F has one more real zero $d \in (33/10, 34/10)$. In fact, $d \approx 3.32178369$. This F gives our desired example, see Figure 1.

(ii) Consider $F(x) = 4x^2 + \pi^2(\cos(x) - 1)$ that has a double zero at 0 and also vanishes at $\pm\pi/2$. Moreover, $F'(x) = 8x - \pi^2 \sin(x)$ vanishes at $x = 0$, $F''(x) = 8 - \pi^2 \cos(x)$ has no common zeroes with F and, for any $k > 1$, $|F^{(2k)}(x)| = \pi^2 |\cos(x)|$ vanishes at $x = \pi/2$ and $|F^{(2k-1)}(x)| = \pi^2 |\sin(x)|$ vanishes at $x = 0$.

A similar example for $n = 3$ is $F(x) = 4x^3 - 6\pi x^2 + \pi^3(1 - \cos(x))$, that vanishes at $0, \pi$ (double zeroes) and $\pi/2$.

Acknowledgements

The authors are supported by Ministerio de Ciencia, Innovación y Universidades of the Spanish Government through grants MTM2016-77278-P (MINECO/AEI/FEDER, UE, first and second authors) and (MTM2017-86795-C3-1-P third author). The three authors are also supported by the grant 2017-SGR-1617 from AGAUR, Generalitat de Catalunya.

References

- [1] E. Casas-Alvero. *Higher order polar germs*. J. Algebra 240 (2001), 326–337.
- [2] Interview to E. Casas-Alvero in Spanish. <https://www.gaussianos.com/la-conjetura-de-casas-alvero-contada-por-eduardo-casas-alvero/>
- [3] W. Castryck, R. Laterveer, M. Ounaïes. *Constraints on counterexamples to the Casas-Alvero conjecture and a verification in degree 12*. Mathematics of Computation 83 (2014), 3017–3037.
- [4] M. Chellali. *On the number of real polynomials of the Casas-Alvero type*. J. of Taibah Univ. for Science 9 (2015), 351–356.
- [5] G. M. Díaz-Toca, L. González-Vega. *On analyzing a conjecture about univariate polynomials and their roots by using Maple*. A Maple conference 2006. Proc. of the conference, Waterloo, Ontario, Canada, July 23–26, 2006. Waterloo: Maplesoft, 2006, p. 81–98.

- [6] J. Draisma, J. P. de Jong. *On the Casas-Alvero conjecture*. Eur. Math. Soc. Newsl. 80 (2011), 29–33.
- [7] H.-C. Graf von Bothmer, O. Labs, J. Schicho, C. van de Woestijne. *The Casas-Alvero conjecture for infinitely many degrees*. J. Algebra 316 (2007), 224–230.
- [8] S. Yakubovich, *Polynomial problems of the Casas-Alvero type*. J. Class. Anal. 4 (2014), 97–120.