

\mathcal{C}^1 SELF-MAPS ON SOME COMPACT MANIFOLDS WITH INFINITELY MANY HYPERBOLIC PERIODIC ORBITS

JUAN LUIS GARCÍA GUIRAO¹, JAUME LLIBRE² AND WEI GAO³

ABSTRACT. The aim of the present work is to provide sufficient conditions for having infinitely many periodic points for \mathcal{C}^1 self-maps having all their periodic orbits hyperbolic and defined on a compact manifold without boundary. The tool used for proving our results is the Lefschetz fixed point theory.

1. INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

Let \mathbb{M} be a topological space and $f : \mathbb{M} \rightarrow \mathbb{M}$ be a continuous map. A point x is called *fixed* if $f(x) = x$ and *periodic* of period k if $f^k(x) = x$ and $f^i(x) \neq x$ if $1 \leq i < k$. By $\text{Per}(f)$ we denote the set of periods of all the periodic points of f .

If $x \in \mathbb{M}$ the set $\{x, f(x), f^2(x), \dots, f^n(x), \dots\}$ is called the *orbit* of the point x . Here f^n means the composition of n times of f with itself. To study the dynamics of the map f is to study all the different kind of orbits of f . Of course, if x is a periodic point of f of period k , then its orbit is $\{x, f(x), f^2(x), \dots, f^{k-1}(x)\}$, and it is called a *periodic orbit*.

In this paper we study the periodic structure of \mathcal{C}^1 self-maps f defined on a given compact manifold \mathbb{M} without boundary. Often the periodic orbits play an important role in the general dynamics of a map, for studying them we can use topological information. Perhaps the best known example in this direction are the results contained in the seminal paper entitle *Period three implies chaos* for continuous self-maps on the interval, see [11].

For continuous self-maps on compact manifolds one of the most useful tools for proving the existence of fixed points and in general of periodic points, is the *Lefschetz Fixed Point Theorem* and its improvements, see for instance [1, 2, 3, 4, 6, 7, 9, 13, 14]. The *Lefschetz zeta function* $\mathcal{Z}_f(t)$ simplifies the study of the periodic points of f . This is a generating function for the Lefschetz numbers of all iterates of f , see section 2 for a precise definition and results about it.

Key words and phrases. Hyperbolic periodic point, period, Lefschetz zeta function, Lefschetz number, sphere.

2010 Mathematics Subject Classification: 58F20.