

LIMIT CYCLES IN PIECEWISE POLYNOMIAL SYSTEMS ALLOWING A NON-REGULAR SWITCHING BOUNDARY

TAO LI¹ AND JAUME LLIBRE²

ABSTRACT. Continuing the investigation for the piecewise polynomial perturbations of the linear center $\dot{x} = -y, \dot{y} = x$ from [Physica D **371**(2018), 28-47] for the case where the switching boundary is a straight line, in this paper we allow that the switching boundary is non-regular, i.e. we consider a switching boundary which separates the plane into two angular sectors with angles $\alpha \in (0, \pi]$ and $2\pi - \alpha$. Moreover, unlike the aforementioned work, we allow that the polynomial differential systems in the two sectors have different degrees. Depending on α and for arbitrary given degrees we provide an upper bound for the maximum number of limit cycles that bifurcate from the periodic annulus of the linear center using the averaging method up to order N . The reachability of the upper bound is also reached for the first two orders. On the other hand, we pay attention to the perturbation of the linear center inside this class of all piecewise polynomial Liénard systems and give some better upper bounds in comparison with the one obtained in the general piecewise polynomial perturbations. Again our results imply that the non-regular switching boundary (i.e. when $\alpha \neq \pi$) the piecewise polynomial perturbations usually leads to more limit cycles than the regular case (i.e. when $\alpha = \pi$) where the switching boundary is a straight line.

1. INTRODUCTION AND STATEMENT OF MAIN RESULTS

In the qualitative theory of smooth differential systems, a classical and challenging objective is to determine the maximum number of the limit cycles bifurcating from the periodic annulus of the linear center $\dot{x} = -y, \dot{y} = -x$, when it is perturbed inside the family formed by all planar polynomial differential systems of the form

$$(1) \quad (\dot{x}, \dot{y}) = \left(-y + \sum_{i=1}^N \varepsilon^i f_i(x, y), x + \sum_{i=1}^N \varepsilon^i g_i(x, y) \right),$$

where $|\varepsilon| > 0$ sufficiently small, and f_i and g_i are real polynomials of degree n . This is essentially the weak Hilbert's 16th problem, see [1, 14, 18]. It was proved in [14] that system (1) has at most $[N(n-1)/2]$ limit cycles bifurcating from the periodic annulus for $|\varepsilon| > 0$ sufficiently small, where as usual $[\cdot]$ denotes the integer part function. Since this upper bound obtained in [14] is not reached in general, up to now we still do not know what is the exact maximum number of limit cycles under the general polynomial perturbation (1) except some special families of perturbations, such as the Liénard family, i.e. $g_i(x, y) = 0$ and $f_i(x, y)$ is independent of the variable y , for which it was proved in [12] that at most $[(n-1)/2]$ limit cycles bifurcate and this number is reached due to [20].

2010 *Mathematics Subject Classification.* 34C29, 34C25, 34C05.

Key words and phrases. Averaging method, Hilbert's 16th problem, limit cycles, discontinuous piecewise polynomial systems, Liénard systems.