

# Minimal set of periods for continuous self–maps of a bouquet of circles

Jaume Llibre

*Departament de Matemàtiques, Universitat Autònoma de Barcelona, Bellaterra  
Barcelona, 08193, Spain  
jllibre@mat.uab.cat*

Ana Sá

*Departamento de Matemática, Faculdade de Ciências e Tecnologia,  
Universidade Nova de Lisboa  
2829-516 Caparica, Portugal  
ams@fct.unl.pt*

Received (to be inserted by publisher)

Let  $G_k$  be a bouquet of circles; i.e. the quotient space of the interval  $[0, k]$  obtained by identifying all points of integer coordinates to a single point, called the branching point of  $G_k$ . Thus,  $G_1$  is the circle,  $G_2$  is the eight space and  $G_3$  is the trefoil. Let  $f : G_k \rightarrow G_k$  a continuous map such that for  $k > 1$  the branching point is fixed.

If  $\text{Per}(f)$  denotes the set of periods of  $f$ , the minimal set of periods of  $f$ , denoted by  $\text{MPer}(f)$ , is defined as  $\bigcap_{g \simeq f} \text{Per}(g)$  where  $g : G_k \rightarrow G_k$  is homological to  $f$ .

The sets  $\text{MPer}(f)$  are well-known for circle maps. Here, we classify all the sets  $\text{MPer}(f)$  for self-maps of the eight space.

*Keywords:* periods, periodic orbits, eight space, continuous maps, set of periods.

## 1. Introduction and statement of the results

In dynamical systems it is often the case that topological information can be used to study qualitative or quantitative properties of the system. This work deals with the problem of determining the set of periods of the periodic orbits of a map given the homology class of the map.

A *finite graph* (simply a *graph*)  $G$  is a topological space formed by a finite set of points  $V$  (points of  $V$  are called *vertices*) and a finite set of open arcs (called *edges*) in such a way that each open arc is attached by its endpoints to vertices. An open arc is a subset of  $G$  homeomorphic to the open interval  $(0, 1)$ . Note that a finite graph is compact, since it is the union of a finite number of compact subsets (the closed edges and the vertices). Notice that a closed edge is homeomorphic either to the closed interval  $[0, 1]$ , or to the circle. It may be either connected or disconnected, and it may have isolated vertices.

The *valence* of a vertex is the number of edges with the vertex as an endpoint (where the closed edges homeomorphic to a circle are counted twice). The vertices with valence 1 of a connected graph are *endpoints* of the graph and the vertices with valence larger than 2 are *branching points*.

Suppose that  $f : G \rightarrow G$  is a continuous map, in what follows a *graph map*. A *fixed point* of  $f$  is a point  $x$  in  $G$  such that  $f(x) = x$ . We will call  $x$  a *periodic point of period  $n$*  if  $x$  is a fixed point of  $f^n$  but it is not fixed by any  $f^k$  for  $1 \leq k < n$ . We denote by  $\text{Per}(f)$  the set of natural numbers corresponding to