

POLYNOMIAL AND RATIONAL FIRST INTEGRALS FOR NON–AUTONOMOUS POLYNOMIAL HAMILTONIAN SYSTEMS

AZUCENA CAICEDO¹, JAUME LLIBRE², AND CLAUDIO VIDAL³

¹Departamento de Matemática, Facultad de Ciencias, Universidad del Bío-Bío,
Casilla 5-C, Concepción, VIII-Región, Chile. e-mail: azu_bac@hotmail.com

²Departament de Matemàtiques, Universitat Autònoma de Barcelona, 08193
Bellaterra, Barcelona, Catalonia, Spain. e-mail: jllibre@mat.uab.cat

³Grupo de Investigación en Sistemas Dinámicos y Aplicaciones-GISDA,
Departamento de Matemática, Facultad de Ciencias, Universidad del Bío-Bío,
Casilla 5-C, Concepción, VIII-Región, Chile. e-mail: clvidal@ubiobio.cl

ABSTRACT. Known results on the existence of polynomial and rational first integrals for autonomous polynomial Hamiltonian systems are extended to non–autonomous polynomial Hamiltonian systems invariant under an involution.

The key tool for proving these results is the existence of Darboux polynomials for the non–autonomous polynomial Hamiltonian systems.

AMS (MOS) Subject Classification. 37J35, 37K10.

1. Introduction and statement of the main results

In this work we deal with the non–autonomous polynomial Hamiltonian systems

$$(1.1) \quad \dot{q}_i = \frac{\partial H(q, p, t)}{\partial p_i}, \quad \dot{p}_i = -\frac{\partial H(q, p, t)}{\partial q_i}, \quad i = 1, \dots, m,$$

with Hamiltonian $H(q, p, t)$, where $q = (q_1, \dots, q_m) \in \mathbb{C}^m$ and $p = (p_1, \dots, p_m) \in \mathbb{C}^m$ are the generalized coordinates. The dot denotes derivative with respect to the time t . Usually q is called the *position vector* and p the *momenta vector*.

In this work we extend the results on the autonomous polynomial Hamiltonian differential systems obtained in [9] to the non–autonomous polynomial Hamiltonian systems (1.1). More precisely, using the existence of some involution under which the Hamiltonian system (1.1) remains invariant and the existence of Darboux polynomials, we provide sufficient conditions in order that the non–autonomous polynomial Hamiltonian systems (1.1) have a second polynomial or rational first integral independent of the Hamiltonian first integral $H(q, p, t)$. These first integrals are polynomial or rational in the variables q and p with coefficients C^1 functions in the time $t \in \mathbb{R}$.