

SOLVING POLYNOMIALS WITH ORDINARY DIFFERENTIAL EQUATIONS

ARMENGOL GASULL AND HECTOR GIACOMINI

ABSTRACT. In this work we consider a given root of a family of n -degree polynomials as a one-variable function that depends only on the independent term. Then we prove that this function satisfies several ordinary differential equations (ODE). More concretely, it satisfies several simple separated variables ODE, a first order generalized Abel ODE of degree $n - 1$ and an $(n - 1)$ -th order linear ODE. Although some of our results are not new, our approach is simple and self-contained. For $n = 2, 3$ and 4 we recover, from these ODE, the classical formulas for solving these polynomials.

1. INTRODUCTION AND MAIN RESULTS

It is known that although general polynomial equations of degree $n \geq 5$ can not be solved by radicals, their roots can be obtained in terms of elliptic or hyperelliptic functions, their inverses or other trascendental functions, like hypergeometric or theta functions. This is a classical subject which starts with results of Hermite, Kronecker and Brioschi and continues with contributions of many others authors, see for instance [1, 7, 9, 12] and the references therein. We will not try to survey all the different points of view from which the question of solving polynomials is addressed.

Because we usually work on ordinary differential equations (ODE) we simply decided to explore which kind of results about polynomial equations can be obtained by using ODE as a main tool. As we will see, our results are self-contained and recover some of the known results on the subject. Before we state our contributions and compare them with these known results, we present a brief survey of the most relevant results that we have found on this subject that also use ODE as a main tool. To the best of our knowledge this approach started with the contributions of Betti ([2]) in 1854 and the ones around 1860 of Cockle and Hartley ([4, 10]). In fact, Enrico Betti proved that the solutions of general polynomial equations satisfy a separated variables ODE and using this fact that he proved that the solutions of these equations can be obtained in terms of hyperelliptic functions and their inverses. He also proved that for quintic equations it suffices to consider elliptic functions and their inverses. On the other hand, James Cockle and Robert Harley showed explicit linear ODE satisfied for a solution of an arbitrary trinomial polynomial equation in terms of its coefficients. For instance, they found a linear homogeneous ODE of 4-th order for a solution $x(q)$ of the quintic polynomial equation in the Bring-Jerrard form $x^5 - x + q = 0$. These results are presented and extended a little in the 1865 Boole's book [3, pp. 190–199]. In his Thesis (“première thèse”), published as a book

2010 *Mathematics Subject Classification*. Primary: 26C05; Secondary: 12D10; 12E12; 13P15; 33C75; 33E05.

Key words and phrases. Polynomial equation; Ordinary differential equation; Abel equations; Elliptic and hyperelliptic integrals.

in 1874 and as an article in 1875 ([14]), Tannery consider a more general question and proved that each branch $y = y(x)$ of an algebraic curve $F(x, y) = 0$, of degree n in y satisfies an n -th order linear homogeneous ODE. In 1887, Heymann ([8]) showed that a solution of a trinomial equation with only one parameter satisfies a linear ODE and realizes for the first time that this solution can be expressed as a hypergeometric function. Mellin ([11]) around 1915 published the quoted paper and others about the representation of the solutions of polynomial equations in terms of multiple integrals, proving again that the solution of a trinomial equations satisfy a linear ODE. Owing to the complexity of explicitly constructing these ODE for non-trinomial polynomial equations, some authors decided to use also functions of several variables, and their corresponding partial differential equations, to express the roots of arbitrary polynomial equations. Bellardinelli, in his extensive review work of 1960 ([1]) presents this type of results and discuss also the previous works about ODE.

We want to stress that results like the one of Tannery have practical applications when people deal with some generating functions appearing in combinatorial problems, see for instance the nice paper of Comtet([5]) and his classical book on Combinatorics ([6]). There, the author gives also a proof that the branches of algebraic curves satisfy linear ODE and several applications of this fact.

In a few words, in this paper we will recover with independent proofs some of the above results and give a few new ones. More concretely, we will re-obtain the results of Betti as a particular case of our more general result and we will give a simple and constructive procedure to obtain non homogeneous $(n-1)$ -order linear ODE satisfied by the solutions of general polynomial equations of degree n . During the way, we prove that these solutions satisfy also Abel type polynomial differential equations of degree $n-1$. We also apply all our results for small n and to some particular examples. It is funny to observe for instance that we obtain the celebrated Cardano's formula for $n=3$, by reducing the computations to solve the simple second order ODE of the harmonic oscillator $x'' = -x$.

Before stating our main results, we start giving an idea of our approach to the problem. In principle, we do not consider all real polynomials, but generic ones. Let $R(x)$ be a monic degree n real polynomial satisfying $R(0) = 0$ and $R'(0) \neq 0$. We are interested in the degree n polynomial equation

$$P(x) = R(x) - q = 0, \quad (1)$$

$q \in \mathbb{R}$, and more in particular, in finding a local explicit expression of the analytic solution $x(q)$ of (1) such that $\lim_{q \rightarrow 0} x(q) = 0$, which exists and is unique by the implicit function theorem because $R'(0) \neq 0$.

Let F be any invertible diffeomorphism such that $F(0) = 0$. Then, from (1) we have the locally equivalent equation $F(R(x)) = F(q)$. We know that $F(R(x(q))) = F(q)$. By derivating it with respect to q , and multiplying both sides by $G(R(x(q))) = G(q)$, with G an arbitrary continuous function, we obtain an ODE with separated variables for $x(q)$

$$G(R(x(q)))F'(R(x(q)))R'(x(q))x'(q) = G(q)F'(q),$$

with initial condition $x(0) = 0$. It can be easily solved giving rise to

$$\phi(x) := \int_0^x G(R(s))F'(R(s))R'(s) ds = \int_0^q G(t)F'(t) dt =: \varphi(q). \quad (2)$$

Hopefully, this new equation $\phi(x) = \varphi(q)$, which is equivalent to $R(x) = q$, allows to obtain $x = \phi^{-1}(\varphi(q))$ being $\phi^{-1} \circ \varphi$ an explicit computable function that it is not trivially equivalent to $x = R^{-1}(q)$. A key point is to chose a suitable F that provides **some cancellation** in the expression $F'(R(x))R'(x)$.

As we will see, this wished cancellation can be obtained from the next proposition, where $\text{dis}_x(P(x))$ denotes the discriminant of $P(x)$ with respect to x . Recall that if $P_n(x) = a_n x^n + \dots + a_1 + a_0$, $a_n \neq 0$,

$$\text{dis}_x(P(x)) = \frac{(-1)^{\frac{n(n-1)}{2}}}{a_n} \text{Res}_x(P_n(x), P_n'(x)),$$

where $\text{Res}_x(P_n(x), Q_m(x))$ is the resultant between P_n and Q_m and can be computed as the $n + m$ determinant

$$\text{Res}_x(P_n(x), Q_m(x)) = \begin{vmatrix} a_n & a_{n-1} & a_{n-2} & \cdots & a_1 & a_0 & 0 & 0 & 0 \\ 0 & a_n & a_{n-1} & a_{n-2} & \cdots & a_1 & a_0 & 0 & 0 \\ 0 & 0 & \ddots & \ddots & \ddots & \cdots & \ddots & \ddots & 0 \\ 0 & 0 & 0 & a_n & a_{n-1} & a_{n-2} & \cdots & a_1 & a_0 \\ b_m & b_{m-1} & \cdots & b_1 & b_0 & 0 & 0 & 0 & 0 \\ 0 & b_m & b_{m-1} & \cdots & b_1 & b_0 & 0 & 0 & 0 \\ 0 & 0 & \ddots & \ddots & \cdots & \ddots & \ddots & \ddots & 0 \\ 0 & 0 & 0 & b_m & b_{m-1} & \cdots & b_1 & b_0 & 0 \\ 0 & 0 & 0 & 0 & b_m & b_{m-1} & \cdots & b_1 & b_0 \end{vmatrix}$$

where $Q_m(x) = b_m x^m + \dots + b_1 x + b_0$, $b_m \neq 0$. It turns out that P_n and Q_m share some (real or complex) root if and only if this determinant vanishes. Hence P_n has some multiple root if and only if its discriminant vanishes, see [13].

Proposition 1.1. *Let $P(x) = R(x) - q$ be a real polynomial of degree $n \geq 2$, with $R(0) = 0$ and $q \in \mathbb{R}$. Set $D(q) = \text{dis}_x(P(x))$. Then*

$$D(R(x)) = (R'(x))^2 U(x) = (P'(x))^2 U(x), \quad (3)$$

for some polynomial U of degree $(n-1)(n-2)$. Moreover, if all the roots of R are simple, $U(0) \neq 0$.

Next theorem is our first main result and proves that a root of a generic polynomial equations $P(x) = R(x) - q = 0$, for q in a neighborhood of 0, can be obtained in terms of hyperelliptic functions and their inverses. As we will comment in Remark 2.1, the restriction that all the roots of R are simple can be removed obtaining a similar result. As we have commented, this result is similar, but more general, to the one given by Betti.

Theorem 1.2. *Let $P(x) = R(x) - q$ be a real polynomial of degree $n \geq 2$, with $R(0) = 0$ and $q \in \mathbb{R}$. Set $D(q) = \text{dis}_x(P(x))$ and assume that all the roots of R are simple. Define the polynomials $\mathcal{D}(q) = \text{sgn}(D(0))D(q)$ and $\mathcal{U}(x) = \mathcal{D}(R(x))/(R'(x))^2$, and the functions*

$$\phi(x) = \text{sgn}(R'(0)) \int_0^x \frac{G(R(s))}{\sqrt{\mathcal{U}(s)}} ds \quad \text{and} \quad \varphi(q) = \int_0^q \frac{G(t)}{\sqrt{\mathcal{D}(t)}} dt, \quad (4)$$

where G is any continuous function satisfying $G(0) \neq 0$. Then, in a neighborhood of 0, ϕ is invertible and

$$x = \phi^{-1}(\varphi(q))$$

is a root of $P(x) = 0$ that goes to 0 as q tends to 0.

In particular, if G is polynomial, ϕ and φ are elliptic or hyperelliptic integrals.

As an illustration, we apply our results to the low degree cases. In particular, when $n = 2$ and $n = 3$ we reobtain the Babylonian and Cardano's formulas, see Sections 2.1 and 2.2, respectively. In Section 2.3 we apply them to the quartic case. Finally, in Section 2.4 we reproduce the results of Betti's work for quintic equations with our point of view.

To state our second main result we recall some definitions. Given $0 < m \in \mathbb{N}$ we will say that a non autonomous first order real ODE of the form

$$x' = a_m(q)x^m + a_{m-1}(q)x^{m-1} + \cdots + a_2(q)x^2 + a_1(q)x + a_0(q) \quad (5)$$

is a *generalized Abel ODE of degree m* . Notice that for $m = 1, 2$ and 3 these equations are usually called *linear*, *Riccati* and *Abel ODE*, respectively. All of them are a subject of classical interest in mathematics.

By using Corollary 2.3, we prove:

Theorem 1.3. *Let $P(x) = R(x) - q$ be a real polynomial of degree $n \geq 2$, with $R(0) = 0$ and $q \in \mathbb{R}$. Let $x(q)$ be one of the roots of this equation, defined in a neighborhood of 0, that tends to zero as q tends to 0. Then $x(q)$ satisfies a generalized Abel ODE (5) of degree $m = n - 1$, where $a_j(q), j = 0, 1, \dots, n - 1$ are rational functions with coefficients depending on the coefficients of R .*

A straightforward consequence of this result is the following corollary.

Corollary 1.4. *Let $P(x) = R(x) - q$ be a real quadratic, cubic or quartic polynomial equation with $R(0) = 0$. Let $x(q)$ be one of the roots of this equation, defined in a neighborhood of 0, that tends to zero as q tends to 0. Then $x(q)$ satisfies, respectively, a linear, Riccati or Abel ODE whose coefficients are rational functions in q .*

The proof of the above results, together with the explicit ODE when $n \in \{2, 3, 4\}$ and P has the canonical form $P(x) = x^n + px - q$ are given in Section 3.

A second consequence of the above results is:

Theorem 1.5. *Let $P(x) = R(x) - q$ be a real polynomial of degree $n \geq 2$, with $R(0) = 0$ and $q \in \mathbb{R}$. Let $x(q)$ be one of the roots of this equation, defined in a neighborhood of 0, that tends to zero as q tends to 0. Then $x = x(q)$ satisfies a $(n - 1)$ -th order linear ODE,*

$$b_{n-1}(q)x^{(n-1)} + b_{n-2}(q)x^{(n-2)} + \cdots + b_1(q)x' + b_0(q)x + b_n(q) = 0,$$

where the functions $b_j(q)$ are polynomials in q , with coefficients depending on the coefficients of R .

Our proof provides a constructive algorithm to obtain all the functions b_j . As we will see in Section 4, when the equation $P(x) = 0$ is the trinomial one, $P(x) = x^n + px - q = 0$, these b_j are extremely simple. We obtain them for $n \leq 6$. We also recover again Cardano's formula by showing that for $n = 3$, and with suitable changes of variables, this differential equation can be written as the equation for the harmonic oscillator. For $n = 4$ in Section 4.1 we present three different expressions of its solution $x(q)$, two of them in terms of hypergeometric functions, and also the classical one.

Finally, in a short Appendix, we present some classical ways to solve the cubic and the quartic equations. This is done, not only for completeness, but for obtaining

a simple and suitable way (for our interests) of presenting the solution of the quartic equation.

2. PROOF OF THEOREM 1.2 AND SOME APPLICATIONS

We start proving Proposition 1.1.

Proof of Proposition 1.1. The polynomial $P' = R'$ has degree $n - 1$. We give the proof when all its roots in \mathbb{C} , $\alpha_1, \alpha_2, \dots, \alpha_{n-1}$, are different and moreover $R(\alpha_j) \neq R(\alpha_k)$ unless $j = k$. The general result follows from this generic case.

Consider the system

$$\begin{cases} P(x) = R(x) - q = 0, \\ P'(x) = R'(x) = 0. \end{cases} \quad (6)$$

Recall, that modulus some non-zero constant, the discriminant between P and P' is the resultant. Hence, by the properties of the resultant we know that $D(q)$ is a polynomial of degree $n - 1$ and moreover it vanishes for all values of q for which the above system is compatible.

Thus, system (6) is compatible if and only if $q = R(\alpha_j)$, $j = 1, 2, \dots, n - 1$. Hence

$$D(q) = K(q - R(\alpha_1))(q - R(\alpha_2)) \cdots (q - R(\alpha_{n-1})), \quad K \neq 0.$$

Consider the new polynomial $Q(x) = D(R(x))$, of degree $n(n - 1)$. It is clear that $Q(\alpha_j) = D(R(\alpha_j)) = 0$. Moreover, $Q'(x) = D'(R(x))R'(x)$. Therefore, $Q'(\alpha_j) = D'(R(\alpha_j))R'(\alpha_j) = 0$. As a consequence, all α_j are double roots of $Q(x)$ and (3) holds.

Finally, since $D(0) = C \operatorname{Res}_x(R(x), R'(x))$, with $C \neq 0$, the hypothesis that all the roots of R are simple is equivalent to $D(0) \neq 0$. Since $R'(0) \neq 0$, we get that $U(0) \neq 0$, as we wanted to prove. \square

Proof of Theorem 1.2. Our proof starts with the discussion given in the introduction of the paper and uses the notations introduced there. Recall that (2) writes as

$$\phi(x) = \int_0^x G(R(s))F'(R(s))R'(s) ds = \int_0^q G(t)F'(t) dt = \varphi(q). \quad (7)$$

We take

$$F(t) = \int_0^t \frac{1}{\sqrt{\mathcal{D}(s)}} ds,$$

that is a local diffeomorphism at 0 because $\mathcal{D}(0) = \operatorname{sgn}(D(0))D(0) > 0$ and, as a consequence, $F'(0) = 1/\sqrt{\mathcal{D}(0)} \neq 0$. Then, by Proposition 1.1,

$$F'(R(s)) = \frac{1}{\sqrt{\mathcal{D}(R(s))}} = \frac{1}{\sqrt{(R'(s))^2 \mathcal{U}(s)}} = \frac{\operatorname{sgn}(R'(0))}{R'(s)\sqrt{\mathcal{U}(s)}} \quad (8)$$

and (7) reads as

$$\phi(x) = \operatorname{sgn}(R'(0)) \int_0^x \frac{G(R(s))}{\sqrt{\mathcal{U}(s)}} ds = \int_0^q \frac{G(t)}{\sqrt{\mathcal{D}(t)}} dt = \varphi(q).$$

Notice that $\varphi'(0) = G(0)/\sqrt{\mathcal{D}(0)} \neq 0$. Hence, φ is a local diffeomorphism at 0. The same happens with ϕ , because $\phi'(0) = \operatorname{sgn}(R'(0))G(0)/\sqrt{\mathcal{U}(0)} \neq 0$. Thus $x = \phi^{-1}(\varphi(q))$ as we wanted to prove. \square

Next remark clarifies the situation when some of the hypotheses in Theorem 1.2 are not satisfied.

Remark 2.1. *In Theorem 1.2 the hypothesis that all the roots of R are simple is used to ensure that $R'(0) \neq 0$ and $D(0) \neq 0$. These conditions together with the hypothesis that $G(0) \neq 0$ imply that the functions defined by the hyperelliptic integrals ϕ and φ are invertible at 0. If we do not mind about their invertibility we arrive also to equality $\phi(x) = \varphi(q)$ with these functions given as in (4) of the statement. We also remark that in this situation the value $\text{sgn}(R'(0))$ must be replaced by the sign of $R(s)$ for s in the interval containing 0 and x .*

Moreover, in this situation, if $R'(0) = 0$, that is when near $x = 0$ the polynomial equation $R(x) = q$ writes as $x^k + O(x^{k+1}) = q$, for some $1 < k \in \mathbb{N}$, k smooth branches of solutions, $x_j(q), j = 1, \dots, k$, solve the equation and satisfy $x_j(0) = 0$. This result is a consequence of Weierstrass' preparation theorem. Each one of these branches satisfies the ODE that we are considering and, as a consequence, the equality $\phi(x) = \varphi(q)$ with both functions given in (4). Similar branches appear also when we try to invert ϕ .

We will also need the following corollaries of previous results. Notice that from the first corollary, the functions defined by the hyperelliptic integrals given in Theorem 1.2 are replaced by primitives of rational functions.

Corollary 2.2. *Let $P(x) = R(x) - q$ be a real polynomial of degree $n \geq 2$, with $R(0) = 0$ and $q \in \mathbb{R}$. Set $D(q) = \text{dis}_x(P(x))$ and assume that all the roots of R are simple. Define the polynomial $U(x) = D(R(x))/(R'(x))^2$ and the functions*

$$\Phi(x) = \int_0^x \frac{H(R(s))}{R'(s)U(s)} ds \quad \text{and} \quad \Psi(q) = \int_0^q \frac{H(t)}{D(t)} dt,$$

where H is any continuous function satisfying $H(0) \neq 0$. Then, in a neighborhood of 0, Φ is invertible and

$$x = \Phi^{-1}(\Psi(q))$$

is a root of $P(x) = 0$ that goes to 0 as q tends to 0.

In particular, if H is polynomial, Φ and Ψ are primitives of rational functions.

Proof. To prove this result we take

$$G(t) = \frac{H(t)}{\sqrt{D(t)}}$$

in Theorem 1.2. Notice that by using (8) we obtain that

$$G(R(s)) = \frac{H(R(s))}{\sqrt{D(R(s))}} = \frac{\text{sgn}(R'(0))H(R(s))}{R'(s)\sqrt{U(s)}}.$$

Therefore,

$$\text{sgn}(R'(0)) \frac{G(R(s))}{\sqrt{U(s)}} = (\text{sgn}(R'(0)))^2 \frac{H(R(s))}{R'(s)(\sqrt{U(s)})^2} = \text{sgn}(U(0)) \frac{H(R(s))}{R'(s)U(s)}.$$

Similarly,

$$\frac{G(t)}{\sqrt{D(t)}} = \frac{H(t)}{(\sqrt{D(t)})^2} = \text{sgn}(D(0)) \frac{H(t)}{D(t)} = \text{sgn}(U(0)) \frac{H(t)}{D(t)}.$$

By replacing both expressions in (4) we obtain that $\Phi(x) = \Psi(q)$ and the corollary follows. \square

This second corollary is essentially a version of Remark 2.1 in this situation. Notice that the hypothesis that all the roots of R are simple it is not needed.

Corollary 2.3. *Let $P(x) = R(x) - q$ be a real polynomial of degree $n \geq 2$, with $R(0) = 0$ and $q \in \mathbb{R}$. Set $D(q) = \text{dis}_x(P(x))$. Let $x = x(q)$ be a root of $P(x) = 0$ that goes to 0 as q tends to 0. Then*

$$x' = \frac{R'(x)U(x)}{D(q)},$$

where U is the polynomial $U(x) = D(R(x))/(R'(x))^2$.

Proof. By Weierstrass' Preparation theorem we know that the algebraic curve $P(x) = R(x) - q$ has at most n branches passing by the point $(x, q) = (0, 0)$. Moreover, each of these branches, say $x = x(q)$, satisfies $R(x(q)) = q$. Hence, $R'(x(q))x'(q) = 1$. From Proposition 1.1, it holds that $D(R(x)) = (R'(x))^2U(x)$ and, as a consequence,

$$x' = \frac{1}{R'(x)} = \frac{R'(x)U(x)}{D(R(x))} = \frac{R'(x)U(x)}{D(q)},$$

as desired. \square

We will apply the above results for $n \leq 5$.

2.1. A toy example: the quadratic equation. Consider $P(x) = x^2 + px - q$, with $p \neq 0$. Then $D(q) = \text{dis}_x(P(x)) = p^2 + 4q$, $\mathcal{D}(q) \equiv D(q)$ and $\mathcal{D}(R(x)) = (2x + p)^2$. Then $\mathcal{U} = 1$. Moreover, since $R'(0) = p$, we get from (4), that for $|q| < p^2/4$,

$$\begin{aligned} \phi(x) &= \int_0^x \frac{\text{sgn}(R'(0))}{\sqrt{\mathcal{U}(s)}} ds = \int_0^x \text{sgn}(p) ds = \text{sgn}(p)x, \\ \varphi(q) &= \int_0^q \frac{1}{\sqrt{\mathcal{D}(t)}} dt = \int_0^q \frac{1}{\sqrt{p^2 + 4t}} dt = \frac{1}{2} \sqrt{p^2 + 4t} \Big|_0^q = \frac{\sqrt{p^2 + 4q} - \sqrt{p^2}}{2}. \end{aligned}$$

Then, by Theorem 1.2 we get equation $\phi(x) = \varphi(q)$, that gives the Babylonian formula

$$x = \frac{-p + \text{sgn}(p)\sqrt{p^2 + 4q}}{2}.$$

By using Corollary 2.2 instead of Theorem 1.2 with $H = 1$ we obtain

$$\begin{aligned} \Phi(x) &= \int_0^x \frac{1}{R'(s)U(s)} ds = \int_0^x \frac{1}{2s + p} ds = \frac{1}{2} \log \left(\frac{2x + p}{p} \right), \\ \Psi(q) &= \int_0^q \frac{1}{D(t)} dt = \int_0^q \frac{1}{p^2 + 4t} dt = \frac{1}{4} \log \left(\frac{p^2 + 4q}{p^2} \right). \end{aligned}$$

By using that $\Phi(x) = \Psi(q)$ we obtain again the classical formula.

Finally, notice that although the obtained formula for $x = x(q)$ is valid when $|q| < p^2/4$, their algebraic nature makes it valid for all values of p and q .

2.2. Cubic equations. We find a solution for the cubic polynomial equation

$$P(x) = x^3 + px - q = 0. \quad (9)$$

Notice the minus sign in front of q , in contrast with the usual notation given in (26) utilized in Section 5.1 of the Appendix. We exclude the trivial case $p \neq 0$.

In the notation of Theorem 1.2, $D(q) = -(4p^3 + 27q^2)$ and $\mathcal{D}(q) = \operatorname{sgn}(p)(4p^3 + 27q^2)$. After some computations,

$$\mathcal{D}(R(x)) = \operatorname{sgn}(p)(3x^2 + 4p)(3x^2 + p)^2,$$

and, as a consequence, $\mathcal{U}(x) = \operatorname{sgn}(p)(3x^2 + 4p)$. Hence, taking $G = 1$, equation $\phi(x) = \varphi(q)$ writes as

$$\int_0^x \frac{\operatorname{sgn}(p)}{\sqrt{\operatorname{sgn}(p)(3s^2 + 4p)}} ds = \int_0^q \frac{1}{\sqrt{\operatorname{sgn}(p)(4p^3 + 27t^2)}} dt. \quad (10)$$

It is well-known that

$$\int_0^x \frac{1}{\sqrt{Ay^2 + B}} dy = \begin{cases} \frac{\operatorname{arcsinh}\left(\sqrt{A/B}x\right)}{\sqrt{A}}, & \text{when } A > 0, B > 0, \\ \frac{\operatorname{arcsin}\left(\sqrt{-A/B}x\right)}{\sqrt{-A}}, & \text{when } A < 0, B > 0, \end{cases} \quad (11)$$

where the second equality is only valid for $|x| < \sqrt{-B/A}$. Thus, for instance applying the first one when $p > 0$ in (10) we obtain that

$$\frac{\sqrt{3}}{3} \operatorname{arcsinh}\left(\frac{\sqrt{3}x}{2\sqrt{p}}\right) = \frac{\sqrt{3}}{9} \operatorname{arcsinh}\left(\frac{3\sqrt{3}q}{2p\sqrt{p}}\right),$$

or equivalently,

$$x = \frac{2\sqrt{p}}{\sqrt{3}} \sinh\left(\frac{1}{3} \operatorname{arcsinh}\left(\frac{3\sqrt{3}q}{2p\sqrt{p}}\right)\right). \quad (12)$$

By using that $\operatorname{arcsinh}(z) = \ln(z + \sqrt{z^2 + 1})$ we obtain that

$$\sinh\left(\frac{1}{3} \operatorname{arcsinh}(z)\right) = \frac{1}{2} \left(\sqrt[3]{z + \sqrt{z^2 + 1}} - \frac{1}{\sqrt[3]{z + \sqrt{z^2 + 1}}} \right)$$

and hence, after some computations, from (12) we get

$$x = \sqrt[3]{\frac{q}{2} + \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}} - \frac{p}{3\sqrt[3]{\frac{q}{2} + \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}}}, \quad (13)$$

that is, Cardano's formula for equation (9).

If we consider the case $p < 0$ and perform the same type of computations but using the second equality in (11) we arrive to

$$x = \frac{2\sqrt{-p}}{\sqrt{3}} \sin\left(\frac{1}{3} \operatorname{arcsin}\left(\frac{3\sqrt{3}q}{2p\sqrt{-p}}\right)\right), \quad (14)$$

that is similar to (12), but for $p < 0$ and only valid when $|q| < \sqrt{-4p^3/27}$.

In any case, as for the quadratic equations, the algebraic nature of the formula (13) allows to consider it for all values of p and q .

2.3. Quartic equations. As we will see, it is difficult to recover the classical solution with this approach. We start with a particularly simple case. We will return to this case in Section 4.1.

Consider the particular quartic equation

$$P(x) = x^4 - 2x^3 + 2x^2 - x - q = 0. \quad (15)$$

We will apply Theorem 1.2 with $G = -2$. After some calculations we obtain that

$$\mathcal{U}(x) = (2x^2 - 2x + 1)^2(4x^2 - 4x + 3) \quad \text{and} \quad D(q) = (4q + 1)^2(16q + 3).$$

Hence,

$$\begin{aligned} \phi(x) &= \operatorname{sgn}(R'(0)) \int_0^x \frac{G(R(s))}{\sqrt{\mathcal{U}(s)}} ds = \int_0^x \frac{2}{(2s^2 - 2s + 1)\sqrt{4s^2 - 4s + 3}} ds \\ &= 2 \arctan \left(\frac{2x - 1}{\sqrt{4x^2 - 4x + 3}} \right) + \frac{\pi}{3} \end{aligned}$$

and

$$\varphi(q) = \int_0^q \frac{G(t)}{\sqrt{\mathcal{D}(t)}} dt = \int_0^q \frac{-2}{(4q + 1)\sqrt{16q + 3}} dt = -\arctan \left(\sqrt{16q + 3} \right) + \frac{\pi}{3}.$$

For the sake of shortness we introduce the new variables

$$z = \frac{2x - 1}{\sqrt{4x^2 - 4x + 3}} \quad \text{and} \quad w = \sqrt{16q + 3}.$$

Notice that given z the corresponding values of x can be obtained by solving a quadratic equation. Hence, the equation $\phi(x) = \varphi(q)$ can be written as

$$2 \arctan(z) = -\arctan(w),$$

or, equivalently, $\tan(2 \arctan(z)) = -w$, that gives

$$\frac{2z}{z^2 - 1} = w.$$

Thus, for each w , the corresponding value of z can be obtained again by solving a new quadratic equation $wz^2 - w - 2z = 0$.

In short, solving two quadratic equations the quartic equation (15) can be solved. In fact, this is the particularity of the equation that we have considered and makes its study easier: there is no need to solve any cubic equation to find its roots. Their four solutions are

$$\frac{1}{2} \pm \frac{1}{2} \sqrt{-1 \pm 2\sqrt{1 + 4q}}.$$

Let us explore what gives our approach when we apply it to a general quartic equation. Recall first, that similarly of what happens with cubic equations, the general quartic case can be reduced to

$$x^4 + px - q = 0, \quad (16)$$

for some $p, q \in \mathbb{R}$. In this situation, a translation is not enough to arrive to (16) and the so-called Tschirnhausen transformations must be used.

If we apply Theorem 1.2 with $G = 1$, we obtain that $D(q) = -(27p^4 + 256q^3)$ and $\mathcal{D}(q) = 27p^4 + 256q^3$. Then, some computations give

$$\mathcal{D}(R(x)) = (R'(x))^2 \mathcal{U}(x) = (4x^3 + p)^2 (16x^6 + 40px^3 + 27p^2).$$

Hence,

$$\phi(x) = \int_0^x \frac{\operatorname{sgn}(R'(0))}{\sqrt{\mathcal{U}(s)}} ds = \int_0^x \frac{1}{\sqrt{16s^6 + 40ps^3 + 27p^2}} ds$$

and

$$\varphi(q) = \int_0^q \frac{1}{\sqrt{\mathcal{D}(s)}} ds = \int_0^q \frac{1}{\sqrt{256s^3 + 27p^4}} ds.$$

The above functions can be expressed as an Appell function and a hypergeometric function, respectively. Therefore, this approach gives no satisfactory results in order to obtain the roots of the quartic equation in terms of radicals. We will return to the quartic equation in Section 4.1.

2.4. Quintic equations. In this section, with our approach, we recover the result of Betti ([2]) that asserts that the solution of these equations can be obtained in terms of the inverse of an elliptic integral. Following Betti, it suffices to study the particular quintic equation

$$P(x) = x^5 + 5x^3 - q.$$

We can not apply directly Theorem 1.2 because the above case is not under its hypotheses. In fact $R(x) = x^5 + 5x^3$ and hence $R'(0) = 0$. Moreover, as we will see, we will use $G(x) = 5\sqrt{5}x$ and thus $G(0) = 0$. Therefore two of the hypotheses of the theorem are not satisfied, but instead we will use the extended result explained in Remark 2.1. There we explain that in this more general situation, it holds that $\phi(x) = \varphi(q)$ with ϕ and φ given also in (4).

Following the notation of Theorem 1.2 we have that

$$\mathcal{D}(q) = 5^5 q^2 (q^2 + 108) \quad \text{and} \quad \mathcal{U}(x) = 5^3 x^2 (x^2 + 5)^2 (x^6 + 4x^4 - 8x^2 + 12).$$

Moreover, as it is explained in Remark 2.1, $\operatorname{sign}(R'(0))$ can be replaced by $+1$, because near 0, R' is positive.

Taking $G(x) = 5\sqrt{5}x$, we get that

$$\phi(x) = \int_0^x \frac{s^2}{\sqrt{s^6 + 4s^4 - 8s^2 + 12}} ds \quad \text{and} \quad \varphi(q) = \int_0^q \frac{1}{5\sqrt{t^2 + 108}} dt.$$

Hence, by introducing the new variable $u = s^2$, the equality $\phi(x) = \varphi(q)$ writes as

$$\int_0^{x^2} \frac{u}{\sqrt{u(u^3 + 4u^2 - 8u + 12)}} du = \int_0^q \frac{2}{5\sqrt{t^2 + 108}} dt.$$

This expression is precisely the one obtained in [2] and gives a root of the considered quintic equation in terms of elementary functions and the inverse of an elliptic integral.

3. POLYNOMIALS AND ABEL EQUATIONS: PROOF OF THEOREM 1.3

This section is devoted to prove Theorem 1.3 and its corollary.

By Corollary 2.3, any branch of solutions $x = x(q)$ of $P(x) = R(x) - q = 0$ passing by $(x, q) = (0, 0)$ satisfies the differential equation

$$x' = \frac{R'(x)U(x)}{D(q)}, \quad (17)$$

where $R'(x)U(x)$ is a polynomial in x of degree $(n-1)^2$ and $D(q)$ is a polynomial in q of degree $n-1$. By dividing $R'U$ by P we get that $R'(x)U(x) = P(x)Q(x) + W(x)$, where W is a polynomial in q and x of degree at most $n-1$ in this last variable. That is,

$$W(x) = \sum_{j=0}^{n-1} w_j(q)x^j,$$

where the functions $w_j(q)$ are polynomials in q . Hence, since $P(x(q)) \equiv 0$, when $x = x(q)$ it holds that

$$x' = \frac{R'(x)U(x)}{D(q)} = \frac{P(x)Q(x) + W(x)}{D(q)} = \frac{W(x)}{D(q)} = \sum_{j=0}^{n-1} \frac{w_j(q)}{D(q)} x^j = \sum_{j=0}^{n-1} a_j(q)x^j, \quad (18)$$

as we wanted to prove.

Let us detail the corresponding ODE (18) for $n = 2, 3, 4$.

For the quadratic equation $P(x) = x^2 + px - q = 0$, we have that

$$D(q) = p^2 + 4q \quad \text{and} \quad U(x) = 1.$$

Hence $x(q)$ satisfies (17),

$$x' = \frac{R'(x)U(x)}{D(q)} = \frac{2x + p}{p^2 + 4q} = \frac{2}{p^2 + 4q}x + \frac{p}{p^2 + 4q}, \quad (19)$$

that is already a linear ODE. Its general solution is

$$x(q) = \frac{-p + K\sqrt{p^2 + 4q}}{2}.$$

By imposing the initial condition $x(0) = 0$ we arrive again to the babylonian solution

$$x(q) = \frac{-p + \operatorname{sgn}(p)\sqrt{p^2 + 4q}}{2}.$$

For the cubic equation $P(x) = x^3 + px - q = 0$,

$$D(q) = -(4p^3 + 27q^2) \quad \text{and} \quad U(x) = -(3x^2 + 4p).$$

Since $R'(x)U(x) = -(9x^4 + 15px^2 + 4p^2) = -9xP(x) - (6px^2 + 9qx + 4p^2)$ and $P(x(q)) \equiv 0$, we have that $x = x(q)$ satisfies the Riccati equation

$$x' = \frac{6px^2 + 9qx + 4p^2}{4p^3 + 27q^2} = \frac{6p}{4p^3 + 27q^2}x^2 + \frac{9q}{4p^3 + 27q^2}x + \frac{4p^2}{4p^3 + 27q^2}. \quad (20)$$

Finally, we consider the quartic equation $P(x) = x^4 + px - q = 0$. Here we have

$$D(q) = -(27p^4 + 256q^3) \quad \text{and} \quad U(x) = -(16x^6 + 40px^3 + 27p^2)$$

and

$$\begin{aligned} R'(x)U(x) &= -(64x^9 + 176px^6 + 148p^2x^3 - 27p^3) \\ &= -(64x^5 + 112px^2 + 64qx)P(x) - (36p^2x^3 + 48pqx^2 + 64q^2x + 27p^3). \end{aligned}$$

Hence $x(q)$ satisfies the Abel ODE

$$x' = \frac{36p^2}{27p^4 + 256q^3}x^3 + \frac{48pq}{27p^4 + 256q^3}x^2 + \frac{64q^2}{27p^4 + 256q^3}x + \frac{27p^3}{27p^4 + 256q^3}.$$

4. POLYNOMIALS AND LINEAR ODE: PROOF OF THEOREM 1.5

We prove Theorem 1.5 and we apply it to the low degree cases. We recover again Cardano's formula, obtaining it as a particular solution of the equation for the harmonic oscillator. We also get the solution of quartic equations in terms of a generalized hypergeometric function that gives an alternative expression to the classical algebraic one, also presented in Section 5.2 of the Appendix.

We start proving the following simple lemma.

Lemma 4.1. *Let $x(q)$ be a solution of the polynomial equation $P(x) = R(x) - q = 0$ and consider $v(q) = A(x(q), q)/D^m(q)$, where $0 < m \in \mathbb{N}$, $D(q) = \text{dis}_x(P(x))$ and A is a polynomial. Then $v'(q) = B(x(q), q)/D^{m+1}(q)$, for some new polynomial $B(x, q)$.*

Proof. We have that

$$v'(q) = \frac{\frac{\partial A(x(q), q)}{\partial x}x'(q) + \frac{\partial A(x(q), q)}{\partial q}}{D^m(q)} - m \frac{A(x(q), q)D'(q)}{D^{m+1}(q)} = \frac{B(x(q), q)}{D^{m+1}(q)},$$

where $B(x, q) = \frac{\partial A(x, q)}{\partial x}R'(x)U(x) + \frac{\partial A(x, q)}{\partial q}D(q) - mA(x, q)D'(q)$, we have used Corollary 2.3 and U is the polynomial appearing in its statement. \square

Proof of Theorem 1.5. We start as in the proof of Theorem 1.3, recalling that by Corollary 2.3 it holds that

$$x' = \frac{R'(x)U(x)}{D(q)} =: \frac{c_1(x)}{D(q)},$$

where c_1 is a polynomial in x of degree $(n-1)^2$. Notice that applying re-iteratively Lemma 4.1, defining $v = x^{(k)}$, $k = 2, 3, \dots$ we obtain that

$$x^{(k)} = \frac{C_k(x, q)}{D^k(q)}, \quad 1 < k \in \mathbb{N},$$

where $C_k(x, q)$ are polynomials of increasing degrees in x , defined recursively as

$$C_{k+1}(x, q) = \frac{\partial C_k(x, q)}{\partial x}R'(x)U(x) + \frac{\partial C_k(x, q)}{\partial q}D(q) - kC_k(x, q)D'(q),$$

and $C_1(x, q) = c_1(x)$. As in the proof of Theorem 1.3, we can write

$$C_k(x, q) = Q_k(x, q)P(x) + B_k(x, q), \quad 0 < k \in \mathbb{N},$$

where each B_k is a polynomial in q and x , of degree at most $n-1$ in this last variable. Hence, $x = x(q)$ satisfies

$$x^{(k)} = \frac{B_k(x, q)}{D^k(q)} = \sum_{j=0}^{n-1} b_{k,j}(q)x^j, \quad 0 < k \in \mathbb{N}. \quad (21)$$

For $k = 1$ this ODE is the one of Abel type given in Theorem 1.3.

Let us explain how to obtain a $(n-1)$ -th order linear differential equation by using (21) for $k = 1, 2, \dots, n-1$. In fact, as a first step, we prove that these $n-1$ ODE can be transformed into $n-2$ ODE, where their left hand sides are polynomials of degree one in the variables $x', x'', \dots, x^{(n-1)}$ and with coefficients that are rational functions of q , while their right hand sides continue being polynomials in x but have decreased their degrees to $n-2$.

If at least $n-2$ of the functions $b_{n-1,j}$ $j = 1, 2, \dots, n-1$, identically vanish, we are done. Otherwise, at least two of them, say $b_{n-1,i}$ and $b_{n-1,\ell}$, are not identically zero. Then, by computing $x^{(i)}/b_{n-1,i}(q) - x^{(\ell)}/b_{n-1,\ell}(q)$ we cancel the term x^{n-1} in the corresponding right hand side, obtaining one of the new desired relations. Doing the same procedure with several couples of relations (21) satisfying that $b_{n-1,j} \neq 0$ we obtain the $n-2$ searched relations.

Starting from these new relations, combining them in a similar way, we obtain, for each $m = 3, \dots, n-1$, in each step $n-m$ ODE whose right hand sides have degree $n-m$ in the variable x . The last step of this procedure gives the desired linear differential equation. \square

Although the linear ODE given in Theorem 1.5 can be obtained in general, their expressions are huge. To show some examples we give these ODE for the particular case of trinomial polynomials

$$P(x) = x^n + px - q = 0, \quad (22)$$

when $n = 3, 4, 5, 6$. As we will see, in this case of trinomial polynomials the resulting ODE has a simple expression.

Notice also that it is not difficult to see that for $n > 1$, near $q = 0$, the solution of (22) is

$$x = x(q) = \frac{1}{p}q - \frac{1}{p^{n+1}}q^n + o(q^n).$$

Hence all the $(n-1)$ -th order linear ODE that we will obtain have to be solved with the initial conditions

$$x(0) = 0, \quad x'(0) = \frac{1}{p}, \quad x''(0) = x'''(0) = \dots = x^{(n-2)}(0) = 0. \quad (23)$$

For $n = 3$, we already know from (20) that

$$x' = \frac{6p}{4p^3 + 27q^2}x^2 + \frac{9q}{4p^3 + 27q^2}x + \frac{4p^2}{4p^3 + 27q^2}.$$

By using the procedure detailed in the proof of Theorem 1.5 we obtain that

$$x'' = -\frac{162pq}{(4p^3 + 27q^2)^2}x^2 + \frac{12p^3 - 162q^2}{(4p^3 + 27q^2)^2}x - \frac{108p^2q}{(4p^3 + 27q^2)^2}.$$

Hence, since $27qx' + (4p^3 + 27q^2)x'' = 3x$, we arrive to the ODE

$$(4p^3 + 27q^2)x'' + 27qx' - 3x = 0. \quad (24)$$

For the non trinomial case the associated ODE is non homogeneous in general (see Remark 4.2). Let us solve it. We introduce a new independent variable t , as $q = g(t)$, for some smooth function g invertible at $t = 0$ and such that $g(0) = 0$. Then $y(t) = x(g(t))$ is a solution of a new second order linear ODE in $y(t)$. Straightforward computations give that the coefficient of $y''(t)$ for this new ODE is $(4p^3 + 27g^2(t))/(g'(t))^2$. Hence, to find a new simple ODE we impose that

$$\frac{4p^3 + 27g^2(t)}{(g'(t))^2} = \operatorname{sgn}(p)3.$$

Solving it we obtain that, when $p \neq 0$, one of its solutions satisfying $g(0) = 0$ and $g'(0) \neq 0$, is

$$g(t) = \begin{cases} \frac{2\sqrt{3}}{9}p\sqrt{p} \sinh(3t), & \text{when } p > 0, \\ \frac{2\sqrt{3}}{9}p\sqrt{-p} \sin(3t), & \text{when } p < 0. \end{cases}$$

In fact, taking these g 's we get that (24) is transformed into the simple equation

$$y'' - \operatorname{sgn}(p)y = 0.$$

Hence, the general solution of (24) is

$$x(q) = \begin{cases} C_1 \sinh(g^{-1}(q)) + C_2 \cosh(g^{-1}(q)), & \text{when } p > 0, \\ C_1 \sin(g^{-1}(q)) + C_2 \cos(g^{-1}(q)), & \text{when } p < 0, \end{cases}$$

where C_1 and C_2 are arbitrary constants. By imposing the initial conditions (23) and computing $g^{-1}(q)$ we get

$$x(q) = \begin{cases} \frac{2\sqrt{p}}{\sqrt{3}} \sinh\left(\frac{1}{3} \operatorname{arcsinh}\left(\frac{3\sqrt{3}q}{2p\sqrt{p}}\right)\right), & \text{when } p > 0, \\ \frac{2\sqrt{-p}}{\sqrt{3}} \sin\left(\frac{1}{3} \operatorname{arcsin}\left(\frac{3\sqrt{3}q}{2p\sqrt{-p}}\right)\right), & \text{when } p < 0, \end{cases}$$

where the second equality takes real values only for $|q| < \sqrt{-4p^3/27}$. These expressions coincide with the ones obtained in Section 2.2, see (12) and (14), and lead us again to Cardano's formula.

The ODE obtained for $n = 4, 5$ and 6 can be obtained similarly. We skip the details and we only show the final results.

For $n = 4$,

$$(27p^4 + 256q^3)x''' + 1152q^2x'' + 688qx' - 40x = 0. \quad (25)$$

For $n = 5$,

$$(256p^5 + 3125q^4)x'''' + 31250q^3x''' + 73125q^2x'' + 31875qx' - 1155x = 0.$$

For $n = 6$,

$$(3125p^6 + 46656q^5)x''''' + 816480q^4x'''' + 4153680q^3x''' + 6658200q^2x'' + 2307456qx' - 57456x = 0.$$

Remark 4.2. For general polynomials (not in trinomial form) it can be seen that the linear differential equations given in Theorem 1.5 are no more homogeneous. This is already the case for polynomials of degree 2, see (19). As an example we give it for $x^3 + sx^2 + px - q = 0$. The associated ODE is

$$(4p^3 + 27q^2 + 18pqs - p^2s^2 - 4qs^3)x'' + (27q + 9ps - 2s^3)x' - 3x - s = 0.$$

When $s = 0$, the above ODE reduces to (24).

4.1. Again quartic equations. To solve the quartic we have to find the solution of (25) with the initial conditions given in (23), that is $x(0) = x''(0) = 0$ and $x'(0) = 1/p$. By using Mathematica we arrive to

$$x = x_1(q) = {}_3F_2 \left(\frac{1}{4}, \frac{1}{2}, \frac{3}{4}; \frac{2}{3}, \frac{4}{3}; -\frac{256q^3}{27p^4} \right) \frac{q}{p}$$

and with Maple we obtain

$$x = x_2(q) = {}_2F_1 \left(-\frac{1}{24}, \frac{5}{24}; \frac{2}{3}; -\frac{256q^3}{27p^4} \right) \cdot {}_2F_1 \left(\frac{7}{24}, \frac{13}{24}; \frac{4}{3}; -\frac{256q^3}{27p^4} \right) \frac{q}{p},$$

where ${}_nF_m(\cdot; \cdot; x)$ are the classical hypergeometric functions.

As an other result for the quartic, by a direct substitution it is easy to check that

$$x = x_3(q) = \frac{1}{2w(q)} \left(\sqrt{2pw^3(q) - 1} - 1 \right),$$

where $w = w(q)$ satisfies $-p^2w^6 + 4qw^4 + 1 = 0$ and $w(0) = 1/\sqrt[3]{p}$, solves it. Notice that this can be done algebraically because there is the algebraic relation between $x_3(q)$ and $w(q)$, $(2x_3w + 1)^2 = 2pw^3 - 1$, and $w(q)$ also satisfies a bi-cubic algebraic equation. See Section 5.2, and in particular (29), to understand how we have obtained the expression x_3 .

In particular, by the uniqueness of solutions theorem, it holds that $x_i(q) = x_j(q)$, for all $i, j \in \{1, 2, 3\}$, although it seems not easy to prove these equalities without passing by the differential equation. It is a challenge to extract the expression of the algebraic solution of (25) by using only the associated ODE.

5. APPENDIX

For completeness we include in this appendix some classical approaches to solve cubic and quartic equations. While for the cubic equations there is nothing new, the solutions of the quartic are given in a form that is not the most commonly used, but that is very practical and it is also suitable for our approach to the problem.

5.1. Cubic equations. The cubic polynomial equations were solved during the XVI Century by the Italian school and the protagonists were Scipione del Ferro, Niccolò Fontana (Tartaglia) and Gerolamo Cardano.

As usual, the cubic polynomial equation $y^3 + by^2 + cy + d = 0$, is transformed into the simpler one

$$x^3 + px + q = 0, \tag{26}$$

for some suitable p and q , by introducing the new variable $x = y + \frac{b}{3}$. We also consider that $p \neq 0$, because otherwise its solutions can be trivially found.

We will recall two different well-known ways for solving it. We start with the most classical one. We look for a solution of the form $x = u + v$. Replacing it in (26) we get $u^3 + v^3 + q + (3uv + p)(u + v) = 0$. Now we impose that u and v simultaneously

satisfy $u^3 + v^3 + q = 0$ and $3uv + p = 0$. By isolating v from the second equation and replacing it into the first one we get that $z = u^3$ satisfies $z^2 + qz - p^3/27 = 0$. Solving this second degree equation and using that $x = u - p/(3u)$ we arrive to the celebrated Cardano's formula,

$$x = \sqrt[3]{-\frac{q}{2} + \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}} - \frac{p}{3\sqrt[3]{-\frac{q}{2} + \sqrt{\frac{q^2}{4} + \frac{p^3}{27}}}}.$$

A different approach is due to François Viète (Vieta). His starting point is the trigonometric identity

$$4 \cos^3(\theta) - 3 \cos(\theta) - \cos(3\theta) = 0. \quad (27)$$

When $p < 0$, we perform in (26) the change of variables $x = u \cos(\theta)$, with $u = 2\sqrt{-p/3}$ and multiply the equation by $4/u^3$. We arrive to

$$4 \cos^3(\theta) - 3 \cos(\theta) - \frac{3q}{2p} \sqrt{\frac{-3}{p}} = 0.$$

Hence, by using (27), when $\left| \frac{3q}{2p} \sqrt{\frac{-3}{p}} \right| \leq 1$, the three solutions of the cubic equation can be obtained from

$$x = 2\sqrt{\frac{-p}{3}} \cos\left(\frac{1}{3} \arccos\left(\frac{3q}{2p} \sqrt{\frac{-3}{p}}\right)\right),$$

taking the different values of the arccos function. When the inequality does not hold or $p > 0$, it is possible to consider the extension of the cos function to \mathbb{C} or to use that the cosh function, $\cosh(x) = (\exp(x) + \exp(-x))/2$, also satisfies

$$4 \cosh^3(\theta) - 3 \cosh(\theta) - \cosh(3\theta) = 0$$

and then use similar ideas to obtain the solutions of the cubic equation.

5.2. Quartic equations. The quartic equation was solved by Ludovico Ferrari, only some few years after the solution of the cubic one. Essentially its solution is based on some tricks for completing squares that strongly use the solution of the cubic equation. As we have already commented, we present a simple and practical version of that approach that is also suitable for our interests.

By a translation, any quartic equation can be written as

$$x^4 + cx^2 + dx + e = 0, \quad d \neq 0, \quad (28)$$

where we discard the trivial case $d = 0$, because then the equation can be easily solved. Trying to get complete squares in both sides we write it as

$$x^4 + (c + u^2)x^2 + \frac{(c + u^2)^2}{4} = u^2x^2 - dx - e + \frac{(c + u^2)^2}{4},$$

for some u to be determined. Therefore it is natural to impose that

$$-e + \frac{(c + u^2)^2}{4} = \frac{d^2}{4u^2} \iff Q(u) := u^6 + 2cu^4 + (c^2 - 4e)u^2 - d^2 = 0.$$

Since $d \neq 0$, any value u satisfying the above bi-cubic equation is non-zero. Hence for any such u , equation (28) writes as

$$\left(x^2 + \frac{c + u^2}{2}\right)^2 = \left(ux - \frac{d}{2u}\right)^2.$$

Then the solutions of (28) coincide with the solutions of the two quadratic equations

$$x^2 + \frac{c + u^2}{2} = \pm \left(ux - \frac{d}{2u}\right).$$

By solving them we obtain that the four solutions of (28) are

$$\frac{u}{2} \left(1 \pm \sqrt{-\frac{2d}{u^3} - \frac{2c}{u^2} - 1}\right), \quad -\frac{u}{2} \left(1 \pm \sqrt{\frac{2d}{u^3} - \frac{2c}{u^2} - 1}\right),$$

where u is any solution of $Q(u) = 0$. Taking $w = 1/u$, them can also be written as

$$\frac{1}{2w} \left(1 \pm \sqrt{-2dw^3 - 2cw^2 - 1}\right), \quad -\frac{1}{2w} \left(1 \pm \sqrt{2dw^3 - 2cw^2 - 1}\right),$$

where w is any solution of the bi-cubic equation

$$-d^2w^6 + (c^2 - 4e)w^4 + 2cw^2 + 1 = 0.$$

In particular, for the trinomial quartic equation $x^4 + px - q = 0$, the solution $x(q)$ that tends to 0 when q also goes to zero is

$$x(q) = \frac{1}{2w} \left(\sqrt{2pw^3 - 1} - 1\right), \quad (29)$$

where $w = w(q)$ satisfies $-p^2w^6 + 4qw^4 + 1 = 0$ and $w(0) = 1/\sqrt[3]{p}$.

In fact, a simple a posteriori proof that the solution $x^4 + px - q = 0$ is given in (29) can be done as follows: write (29) as

$$S(x, w) := (2xw + 1)^2 + 1 - 2pw^3 = 0$$

and observe that

$$\text{Res}_w(S(x, w), -p^2w^6 + 4qw^4 + 1) = -4096p^2(x^4 + px - q)^3,$$

where Res_w denotes the resultant with respect to w , see [13]. Hence, when $p \neq 0$, if w satisfies simultaneously $S(x, w) = 0$ and $-p^2w^6 + 4qw^4 + 1 = 0$, the corresponding x is a zero of the quartic polynomial.

Acknowledgements. The first author is partially supported by the Ministry of Science and Innovation–Research Agency of the Spanish Government through grant PID2019-104658GB-I00 and by AGAUR, Generalitat de Catalunya, through grant 2017-SGR-1617.

REFERENCES

- [1] G. Bellardinelli, *Fonctions hyperg om etriques de plusieurs variables et r solution analytique des  quations alg briques g n rales*, M morial des sciences math matiques, fascicule 145, 1960.
- [2] E. Betti, *Un teorema sulla risoluzione analitica delle equazioni algebriche*. *Annali de Scienze mathematiche e fisiche* **V** (1854), 10–17.
- [3] G. Boole, *Treatise on differential  quations*, Supplementary volume, Mac Millan, Cambridge and London, 1865.
- [4] J. Cockle, *On transcendental and algebraic solution*. *Philosophical Magazine* **21** (1861), 379–383.

- [5] L. Comtet, *Calcul pratique des coefficients de Taylor d'une fonction algébrique*. Enseignement Math. **10** (1964), 267–270.
- [6] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, Dordrecht, Reidel, 1974.
- [7] L. Dickson, *Modern Algebraic Theories*. Benj. H. Sanborn and Co., 1926.
- [8] W. Heymann, *Theorie der trinomische Gleichungen*. Math. Ann. **28** (1887), 61.
- [9] R. B. King, *Beyond the quartic equation*. Birkhäuser Boston, Inc., Boston, MA, 1996.
- [10] R. Harley, *On the Transcendental Solution of Algebraic Equations*. Quart. J. Pure Appl. Math. **5** (1862), 337–361.
- [11] H.J. Mellin, *Zur Theorie der trinominchen Gleichungen*. Ann. Ac. Sc. Fenn. **7** (1915).
- [12] V. Prasolov, Y. Solovyev, *Elliptic functions and elliptic integrals*. Translated from the Russian manuscript by D. Leites. Translations of Mathematical Monographs, **170**. American Mathematical Society, Providence, RI, 1997.
- [13] B. Sturmfels, *Solving systems of polynomial equations*, vol. **97** of CBMS Regional Conference Series in Mathematics. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI (2002).
- [14] J. Tannery, *Propriétés des intégrales des équations différentielles à coefficients variables*. Annales scientifiques de l'ENS (1875), 113–182.

DEPARTAMENT DE MATEMÀTIQUES, EDIFICI Cc, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 CERDANYOLA DEL VALLÈS, BARCELONA, SPAIN.

CENTRE DE RECERCA MATEMÀTICA, EDIFICI Cc, CAMPUS DE BELLATERRA, 08193 CERDANYOLA DEL VALLÈS, BARCELONA, SPAIN.

E-mail address: gasull@mat.uab.cat

INSTITUT DENIS POISSON. UNIVERSITÉ DE TOURS, C.N.R.S. UMR 7013. 37200 TOURS, FRANCE.

E-mail address: Hector.Giacomini@lmpt.univ-tours.fr