



Period function of planar turning points

Renato Huzak¹ and David Rojas ²

¹Hasselt University, Campus Diepenbeek, Agoralaan Gebouw D, 3590 Diepenbeek, Belgium

²Departament d'Informàtica, Matemàtica Aplicada i Estadística, Universitat de Girona, Girona 17003, Catalonia, Spain

Received 17 November 2020, appeared 22 March 2021

Communicated by Armengol Gasull

Abstract. This paper is devoted to the study of the period function of planar generic and non-generic turning points. In the generic case (resp. non-generic) a non-degenerate (resp. degenerate) center disappears in the limit $\epsilon \rightarrow 0$, where $\epsilon \geq 0$ is the singular perturbation parameter. We show that, for each $\epsilon > 0$ and $\epsilon \sim 0$, the period function is monotonously increasing (resp. has exactly one minimum). The result is valid in an ϵ -uniform neighborhood of the turning points. We also solve a part of the conjecture about a uniform upper bound for the number of critical periods inside classical Liénard systems of fixed degree, formulated by De Maesschalck and Dumortier in 2007. We use singular perturbation theory and the family blow-up.

Keywords: critical periods, family blow-up, period function, slow-fast systems.


2020 Mathematics Subject Classification: 34E15, 34E17.

1 Introduction

We consider slow-fast polynomial Liénard equations of center type

$$X_{\epsilon, \eta} : \begin{cases} \dot{x} = y - \left(x^{2n} + \sum_{k=1}^l a_k x^{2n+2k} \right), \\ \dot{y} = \epsilon^{2n} \left(-x^{2n-1} + \sum_{k=1}^m b_k x^{2n+2k-1} \right), \end{cases} \quad (1.1)$$

where $l, m, n \geq 1$, $\eta := (a_1, \dots, a_l, b_1, \dots, b_m)$ is kept in a compact set K of \mathbb{R}^{l+m} and $\epsilon \geq 0$ is the singular perturbation parameter kept small. System $X_{\epsilon, \eta}$ is invariant under the symmetry $(x, t) \rightarrow (-x, -t)$ and has a center at the origin for all $\epsilon > 0$, $\epsilon \sim 0$, and for all $\eta \in K$. The center is non-degenerate when $n = 1$ or nilpotent when $n > 1$. In the limit $\epsilon = 0$, we encounter drastic changes in the dynamics of (1.1): the system has a curve of singular points, given by $\{y = x^{2n} + \sum_{k=1}^l a_k x^{2n+2k}\}$, passing through the origin, and horizontal regular orbits (see Figure 1.1).

 Corresponding author. Email: david.rojas@udg.edu