



MATHEMATICAL SCIENCES

Rational first integrals of the Liénard equations: The solution to the Poincaré problem for the Liénard equations

JAUME LLIBRE, CLAUDIO PESSOA & JARNE D. RIBEIRO

Abstract: Poincaré in 1891 asked about the necessary and sufficient conditions in order to characterize when a polynomial differential system in the plane has a rational first integral. Here we solve this question for the class of Liénard differential equations $\ddot{x} + f(x)\dot{x} + x = 0$, being $f(x)$ a polynomial of arbitrary degree. As far as we know it is the first time that all rational first integrals of a relevant class of polynomial differential equations of arbitrary degree has been classified.

Key words: Liénard equation, rational first integral, Poincaré problem, polynomial differential equation.

1 - THE POINCARÉ PROBLEM ON THE RATIONAL FIRST INTEGRALS OF THE POLYNOMIAL DIFFERENTIAL SYSTEMS

A rational function $f(x,y)/g(x,y)$ has degree m if the polynomials $f(x,y)$ and $g(x,y)$ are coprime in the ring $\mathbb{R}[x,y]$, and the maximum of the degrees of $f(x,y)$ and $g(x,y)$ is m .

A polynomial differential system is a differential system of the form

$$\frac{dx}{dt} = \dot{x} = P(x,y), \quad \frac{dy}{dt} = \dot{y} = Q(x,y), \quad (1)$$

where $P(x,y)$ and $Q(x,y)$ are real polynomials in the variables x and y , and t is the independent variable usually called the *time*. The polynomial vector field associated to the polynomial differential system (1) is

$$\mathcal{X} = P(x,y) \frac{\partial}{\partial x} + Q(x,y) \frac{\partial}{\partial y}.$$

Let U be an open subset of \mathbb{R}^2 . Here a *first integral* is a \mathcal{C}^1 non-locally constant function $H : U \rightarrow \mathbb{R}$ such that it is constant on the solutions $(x(t), y(t))$ of the polynomial differential system (1) contained in U , i.e. if $\mathcal{X}(H)|_U \equiv 0$.

If the function H is rational then we say that H is a *rational first integral*.

The problem of providing necessary and sufficient conditions in order that a polynomial differential system in the plane has a rational first integral was stated by Poincaré (1891). This problem