

RESEARCH

Open Access



Minimal set of periods for continuous self-maps of the eight space

Jaume Llibre^{1*} and Ana Sá²

*Correspondence:
jlilibre@mat.uab.es

¹Departament de Matemàtiques,
Universitat Autònoma de Barcelona,
08193 Bellaterra, Barcelona, Spain
Full list of author information is
available at the end of the article

Abstract

Let G_k be a bouquet of circles, i.e., the quotient space of the interval $[0, k]$ obtained by identifying all points of integer coordinates to a single point, called the branching point of G_k . Thus, G_1 is the circle, G_2 is the eight space, and G_3 is the trefoil. Let $f : G_k \rightarrow G_k$ be a continuous map such that, for $k > 1$, the branching point is fixed.

If $\text{Per}(f)$ denotes the set of periods of f , the minimal set of periods of f , denoted by $\text{MPer}(f)$, is defined as $\bigcap_{g \sim f} \text{Per}(g)$ where $g : G_k \rightarrow G_k$ is homological to f .

The sets $\text{MPer}(f)$ are well known for circle maps. Here, we classify all the sets $\text{MPer}(f)$ for self-maps of the eight space.

MSC: 37E15

Keywords: Continuous maps; Periodic orbit; Period; Minimal period; The space in shape of eight

1 Introduction and statement of the results

In dynamical systems it is often the case that topological information can be used to study qualitative or quantitative properties of the system. This work deals with the problem of determining the set of periods of the periodic orbits of a map given the homology class of the map.

A *finite graph* (simply a *graph*) G is a topological space formed by a finite set of points V (points of V are called *vertices*) and a finite set of open arcs (called *edges*) in such a way that each open arc is attached by its endpoints to vertices. An open arc is a subset of G homeomorphic to the open interval $(0, 1)$. Note that a finite graph is compact since it is the union of a finite number of compact subsets (the closed edges and the vertices). Notice that a closed edge is homeomorphic either to the closed interval $[0, 1]$ or to the circle. It may be either connected or disconnected, and it may have isolated vertices.

The *valence* of a vertex is the number of edges with the vertex as an endpoint (where the closed edges homeomorphic to a circle are counted twice). The vertices with valence 1 of a connected graph are *endpoints* of the graph and the vertices with valence larger than 2 are *branching points*.

Suppose that $f : G \rightarrow G$ is a continuous map, in what follows a *graph map*. A *fixed point* of f is a point x in G such that $f(x) = x$. We will call x a *periodic point of period n* if x is a

© The Author(s) 2021. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.