On the reversible quadratic polynomial vector fields on the two dimensional sphere

Weber F. Pereira[1], <u>Claudio Pessoa</u>[2]

[1] *Department of Mathematics, Universidade Estadual Paulista, 15054-000–S. J. Rio Preto, Brazil.*
E-mail: weberf@ibilce.unesp.br   *URL:* http://www.mat.ibilce.unesp.br/personal/weber.html

[2] *Department of Mathematics, Universidade Estadual Paulista, 15054-000–S. J. Rio Preto, Brazil.*
E-mail: pessoa@ibilce.unesp.br   *URL:* http://www.mat.ibilce.unesp.br/personal/claudio_gomes.html

We study a class of quadratic reversible polynomial vector fields on $\mathbb{S}^2$ with $(3, 2)$-type reversibility. We classify all isolated singularities and we prove the nonexistence of limit cycles for this class. Our study provides tools to determine the phase portrait for these vector fields.

References

[1] C.A. Buzzi, M.A. Teixeira and J. Yang, Hopf–zero bifurcations of reversible vector fields, *Nonlinearity*, **14** (2001), 623–638.

[2] H. Giacomini, J. Llibre and M. Viano, On the nonexistence, existence and uniqueness of Limit cycles, *Nonlinearity* **9** (1996), 501–516.

[3] J. Llibre and C. Pessoa, Homogeneous polynomial vector fields of degree 2 on the 2–dimensional sphere, *Extracta Mathematicae* **21**, n^o 2, (2006), 167–190.

[4] J. Llibre and C. Pessoa, Phase portraits for quadratic homogeneous polynomial vector fields on $\mathbb{S}^2$, *Rend. Cir. Mat. Palermo*, **58** (2009), 361–406.

[5] W. F. Pereira and C. Pessoa, A class of reversible quadratic polynomial vector fields on $\mathbb{S}^2$, *J. Math. Anal. Appl.*, **371** (2010), 203–209.